arXiv:1505.06785v4 [math.CV] 25 Jul 2015

EXTREMAL LENGTH FUNCTIONS ARE LOG-PLURISUBHARMONIC

HIDEKI MIYACHI

Abstract. In this paper, we show that the extremal length functions on Teichmüller space are log-plurisubharmonic. As a corollary, we obtain an alternative proof of L.Liu and W.Su's results on the plurisubharmonicity of extremal length functions. We also obtain alternative proofs of S.Krushkal's results that a function defined by the Teichmüller distance from a reference point is plurisubharmonic, and the Teichmüller space is hyperconvex. To show the log-plurisubharmonicity, we give an explicit formula of the Levi form of the extremal length functions in generic case.

Contents

1. Introduction 1
2. Notation 3
3. Coordinates via representations of the odd cohomology 6
4. Family of quadratic differentials with prescribed vertical foliation 11
5. Levi forms and Plurisubharmonicity of subspecies 15
6. Gemetric interpretation of η_v 19
7. Appendix 22
References 25

1. Introduction

1.1. **Results.** A positive function U on a complex manifold N is said to be *log-plurisubharmonic* if $\log U$ is plurisubharmonic. Any log-plurisubharmonic function is plurisubharmonic (cf. §5.2).

Let $\mathcal{T}_{g,m}$ be the Teichmüller space of Riemann surfaces of analytically finite type (g,m) with $2g-2+m>0$. One of the aims of this paper is to give the following theorem, which improves a result by L.Liu and W. Su in [23].

Theorem 1.1 (Log-plurisubharminicity). *Extremal length functions on $\mathcal{T}_{g,m}$ are log-plurisubharmonic.*

Moreover, we will also observe that for generic measured foliations, the log-extremal length functions are real analytic strictly plurisubharmonic functions on $\mathcal{T}_{g,m}$ (cf. Theorem 5.3).

2010 *Mathematics Subject Classification.* 32G15, 30F45, 31C10.

Key words and phrases. Teichmüller theory, Teichmüller distance, Extremal length, Plurisubharmonic functions.

When $3g-3+m=1$, Theorem 1.1 follows from the direct calculation (cf. §7.2). Hence, in the discussion in the later section, we always assume that $3g-3+m\geq 2$. We will prove Theorem 1.1 in §5.2. In fact, Theorem 1.1 is derived from a more stronger property of the extremal length functions than that given in Theorem 1.1 (cf. Theorem 5.2). However, the property in Theorem 1.1 is also important as we see below (cf. Corollaries 1.1 and 1.2). To show Theorem 1.1, we will give an explicit formula of the Levi forms of extremal length functions in generic case. (cf. Theorem 5.1).

The following immediately follows from Theorem 1.1 (cf. [17, Colollary 2.6.9]).

Corollary 1.1 (Log-plurisubharmonicity of positive polynomials)**.** *Let* $F_1,\cdots,F_n$ *be measured foliations. For any polynomial* P *of* n*-variables with positive coefficients, the function*

$$\mathcal{T}_{g,m}\ni x\mapsto P(\mathrm{Ext}_x(F_1),\cdots,\mathrm{Ext}_x(F_n))\in\mathbb{R}$$

is log-plurisubharmonic.

1.2. **Teichmüller distance is plurisubharmonic.** It is known that the least upper semicontinuous majorant of a family of plurisubharmonic funtions is either a constant function $+\infty$ or plurisubharmonic (cf. Theorem 5 in [21, Chapter II, §2]). Kerckhoff's formula (2.2) implies that the Teichmüller distance $d_T(x_0,\cdot)$ from a reference point $x_0\in\mathcal{T}_{g,m}$ coincides with the half of the least upper semicontinuous majorant of the family of log-extremal length functions. Therefore, we obtain an alternative proof of Krushkal's result [20, Corollary 3] as follows.

Corollary 1.2 (Teichmüller distance is plurisubharmonic)**.** *For* $x_0\in\mathcal{T}_{g,m}$*, the distance function*

$$\mathcal{T}_{g,m}\ni x\mapsto d_T(x_0,x)$$

is plurisubharmonic.

1.3. **Convexities for Teichmüller space.** A subset K in a complex manifold N is said to be *disk-convex* in N if for any continuous mapping $g\colon\overline{\mathbb{D}}\to N$, holomorphic in $\mathbb{D}$, $f(\partial\mathbb{D})\subset K$ implies $f(\overline{\mathbb{D}})\subset K$ (cf. [26]). For $\epsilon>0$ and a measured foliation F, we call the set of the form $\{x\in\mathcal{T}_{g,m}\mid \mathrm{Ext}_x(F)<\epsilon\}$ the *ϵ-horoball for F*. From the maximum modulus principle for (pluri)subharmonic functions, we obtain the following.

Corollary 1.3 (Disk convexity)**.** *For* $\epsilon>0$ *and a measured foliation* F*, the* ϵ*-horoball for* F *is disk-convex in* $\mathcal{T}_{g,m}$*. The metric ball with respect to the Teichmüller distance is also disk-convex in* $\mathcal{T}_{g,m}$*.*

About the convexity of metric balls, A. Lenzhen and K. Rafi [22] observed that the metric ball with respect to the Teichmüller distance is quasi-convex. Furthermore, they also showed that the extremal length function is not convex along a Teichmüller geodesic in general. This means that the intersection between a horoball of extremal length and a Teichmüller disk is simply connected, but may not be hyperbolically convex in general.

A complex manifold N is said to be *hyperconvex* (in the sense of Stehlé) if it admits a continuous plurisubharmonic exhaustion function $\rho\colon N\to[-\infty,0)$ in the sense that $\{x\in N\mid\rho(x)<c\}$ is relatively compact in M for every $c<0$ (cf. [32].

See also [28]). Take two essentially complete measured foliations F and G which are transverse in the sense that

$$i(F,H)+i(G,H)>0 \tag{1.1}$$

for all $H \in \mathcal{MF}-\{0\}$. In Theorem 5.2 given later, we will check that

$$\rho\colon \mathcal{T}_{g,m} \ni x \mapsto -\frac{1}{\mathrm{Ext}_x(F)+\mathrm{Ext}_x(G)+1} \tag{1.2}$$

is a real analytic, negative, strictly plurisubharmonic exhaustion function with lower bounds. Thus we obtain the following.

Corollary 1.4 (Hyperconvexitiy)**.** *Teichmüller space is hyperconvex.*

Corollary 1.4 is already given by Krushkal in [19]. Notice that the hyperconvexity of Teichmüller space also follows from the compleness of the Carathéodory distance (cf. [1] and [7]). As an immediate corollary of the hyperconvexity (or Corollary 1.2), by virtue of Oka's theorem, we deduce the following, which is first proven by L.Bers and L.Ehrenpreis (see [2]. See also [7], [23], [31], [36]).

Corollary 1.5 (Holomorphic convexity)**.** *Teichmüller space is a domain of holomorphy.*

2. NOTATION

2.1. **Teichmüller space.** Let $\Sigma_{g,m}$ be a compact orientable surface of genus g with m-disks removed with $2g-2+m>0$. The *Teichmüller space* $\mathcal{T}_{g,m}$ of Riemann surfaces of analytically finite type (g,m) is the set of equivalence classes of pairs (M,f) consisting of a Riemann surface M of analytically finite type (g,m) and an orientation preserving homeomorphism $f\colon \mathrm{Int}(\Sigma_{g,m}) \to M$. Two marked Riemann surfaces (M_1,f_1) and (M_2,f_2) are *equivalent* if there is a conformal mapping $h\colon M_1 \to M_2$ which homotopic to $f_2 \circ f_1^{-1}$.

The Teichmüller space admits a canonical distance d_T, which we call the *Teichmüller distance.* The Teichmüller distance is originally defined as the logarithm of the infimum of the maximal dilatations of quasiconformal mappings respecting the markings (cf. [14, §5]). S. Kerckhoff [16] gave a geometric description of the Teichmüller distance via the extremal length, which we will recall in (2.2) below.

2.2. **Measured foliations.** Let $\mathcal{S}$ be the set of homotopy classes of non-trivial and non-peripheral simple closed curves on $\Sigma_{g,m}$. Let $\mathcal{WS}$ be the set of weighted simple closed curves, that is, the set of formal products $t\alpha$ of non-negative number t and $\alpha \in \mathcal{S}$. The closure $\mathcal{MF}$ of the embedding

$$\mathcal{WS} \ni t\alpha \mapsto [\mathcal{S} \ni \beta \mapsto t\, i(\alpha,\beta)] \in \mathbb{R}_+^{\mathcal{S}}$$

is called the *space of measured foliations* on $\Sigma_{g,m}$. We consider $\mathcal{WS}$ as a subset of $\mathcal{MF}$. We identify $1\cdot\alpha \in \mathcal{WS}$ with $\alpha \in \mathcal{S}$. Any measured foliation is described as a pair consisting of a singular foliation and a transverse measure (cf. [5, Exposé 5]). For instance, $t\alpha \in \mathcal{MF}$ is a foliated annulus with core α which identified with an annulus $[0,t]\times[0,1]/(x,0)\sim(x,1)$ and a transverse measure associated to $|dx|$.

It is known that the geometric intersection number $i(\alpha,\beta)$ for $\alpha,\beta \in \mathcal{S}$ extends continuously to $\mathcal{MF}\times\mathcal{MF}$ (cf. [3]).

If we fix a complete hyperbolic structure on the interior of $\Sigma_{g,m}$, any measured foliation canonically corresponds to a measured lamination. A *measured lamination* is a pair of a geodesic lamination, which called the *support*, and a transverse measure, where a *geodesic lamination* a is a compact set in the interior fo $\Sigma_{g,m}$ which consists of disjoint complete geodesics (cf. [29, §1.7]). A measured foliation is said to be *essentially complete* if the support of the corresponding measured lamination is maximal, that is, the complement consists of ideal triangles or ideal punctured monogon if $(g, m) \neq (1, 1)$, a punctured bigon otherwise (cf. [33, Definition 9.5.1, Propositions 9.5.2 and 9.5.4]). In the interior of $\Sigma_{g,m}$, each singularity of the associated foliation of an essentially complete measured foliation is a three prong singularity. At any puncture, the associated foliation has a one prong singularity (cf. [29, Epilogue]).

2.3. **Extremal length.** Let M be a Riemann surface and let A be a doubly connected domain on M. If A is conformally equivalent to a round annulus $\{1 < |z| < R\}$, we define the *modulus* of A by

$$\mathrm{Mod}(A) = \frac{1}{2\pi}\log R.$$

The *extremal length* of a simple closed curve α on M is defined by

$$\mathrm{Ext}_M(\alpha) = \inf\left\{\frac{1}{\mathrm{Mod}(A)} \mid \text{the core curve of } A \subset M \text{ is homotopic to } \alpha\right\}. \tag{2.1}$$

In [16], Kerckhoff showed that if we define the extremal length of $t\alpha \in \mathcal{WS}$ by

$$\mathrm{Ext}_M(t\alpha) = t^2\mathrm{Ext}_M(\alpha),$$

then the extremal length function Ext_M on $\mathcal{WS}$ extends continuously to $\mathcal{MF}$. For $F \in \mathcal{MF}$, the extremal length function $\mathrm{Ext}_\cdot(F)$ on $\mathcal{T}_{g,m}$ is defined by

$$\mathrm{Ext}_x(F) = \mathrm{Ext}_M(f(F)).$$

It is known that the function

$$\mathcal{T}_{g,m} \times \mathcal{MF} \ni (x, F) \mapsto \mathrm{Ext}_x(F)$$

is continuous. Furthermore, the extremal length function $\mathrm{Ext}_x \colon \mathcal{MF} \to \mathbb{R}$ is a proper function and satisfies the quasiconformal distortion property:

$$\mathrm{Ext}_y(F) \leq e^{2d_T(x,y)}\mathrm{Ext}_x(F)$$

for any $F \in \mathcal{MF}$ and $x, y \in \mathcal{T}_{g,m}$.

2.4. **Kerckhoff's formula and Minsky's inequality.** The Teichmüller distance d_T is expressed with the extremal length, which we call *Kerckhoff's formula*:

$$\begin{aligned} d_T(x_1, x_2) &= \frac{1}{2}\log\sup_{\alpha\in\mathcal{S}}\frac{\mathrm{Ext}_{x_2}(\alpha)}{\mathrm{Ext}_{x_1}(\alpha)} \\ &= \frac{1}{2}\sup_{\alpha\in\mathcal{S}}\left(\log\mathrm{Ext}_{x_2}(\alpha) - \log\mathrm{Ext}_{x_1}(\alpha)\right) \end{aligned} \tag{2.2}$$

for $x_1, x_2 \in \mathcal{T}_{g,m}$ (see [16]). Y. Minsky [27] observed the following inequality, which we recently call *Minsky's inequality*:

$$i(F, G)^2 \leq \mathrm{Ext}_x(F)\,\mathrm{Ext}_x(G) \tag{2.3}$$

for $x \in \mathcal{T}_{g,m}$ and $F, G \in \mathcal{MF}$. If two measured foliation F and G are transverse in the sense of (1.1), we have

$$\mathrm{Ext}_x(F) + \mathrm{Ext}_x(G) \geq \frac{i(F, H)^2 + i(G, H)^2}{\mathrm{Ext}_x(H)}$$

for all H and hence we deduce

$$\mathrm{Ext}_x(F)+\mathrm{Ext}_x(G)\geq e^{2d_T(x_0,x)}\min\{i(F,H)^2+i(G,H)^2\mid \mathrm{Ext}_{x_0}(H)=1\}.$$

This implies that the function $x\mapsto \mathrm{Ext}_x(F)+\mathrm{Ext}_x(G)$ is proper on $\mathcal{T}_{g,m}$.

2.5. **Quadratic differentials.** Let $\mathcal{Q}_{g,m}$ be the space of holomorphic quadratic differentials with finite L^1-norm. Namely, the space $\mathcal{Q}_{g,m}$ is a holomorphic vector bundle over $\mathcal{T}_{g,m}$ and the fiber $\mathcal{Q}_x$ at $x=(M,f)\in\mathcal{T}_{g,m}$ is the space of holomorphic quadratic differentials on M with finite L^1-norm (cf. §2.7). Any $q\in\mathcal{Q}_x$ extends meromorphically to the completion $\overline{M}$ by filling punctures and has (at most) simple poles at punctures. A quadratic differential $q\in\mathcal{Q}_{g,m}$ is said to be *generic* if it satisfies the following two conditions:

(1) q has a simple pole at every puncture.
(2) all zeroes of q are simple.

We can easily check that the set of all generic holomorphic quadratic differentials is an open and dense subset in $\mathcal{Q}_{g,m}$.

2.6. **Hubbard-Masur differentials.** For $q\in\mathcal{Q}_{g,m}$, there is a unique measured foliation F such that

$$i(F,\alpha)=\inf_{\alpha'\sim f(\alpha)}\int_{\alpha'}|\mathrm{Re}\sqrt{q}|.$$

for all $\alpha\in\mathcal{S}$, where $(M,f)=\pi(q)\in\mathcal{T}_{g,m}$. We call F the *vertical foliation* and denote it by $\mathbf{v}(q)$. We call $\mathbf{h}(q)=\mathbf{v}(-q)$ the *horizontal foliation* of q. It is known that for any $F\in\mathcal{MF}$ and $x=(M,f)\in\mathcal{T}_{g,m}$, there is a unique holomorphic quadratic differential $q_{F,x}$ such that $\mathbf{v}(q_{F,x})=F$. We call $q_{F,x}$ the *Hubbard-Masur differential* for F on x (cf. [13]). If F is essentially complete, the Hubbard-Masur differential $q_{F,x}$ is generic for all $x\in\mathcal{T}_{g,m}$, since any Whitehead equivalent measured foliation to F is isotopic to F.

The Hubbard-Masur differential $q_{F,x}=q_{F,x}(z)dz^2$ for F on $x=(M,f)$ satisfies

$$\mathrm{Ext}_x(F)=\|q_{F,x}\|=\int_M|q_{F,x}|=\int_M|q_{F,x}(z)|\cdot\frac{i}{2}dz\wedge d\overline{z}. \tag{2.4}$$

2.7. **Infinitesimal theory.** Teichmüller space also has a canonical complex structure, which induced from the deformation by quasiconformal mappings. A *Beltrami differential* is, by definition, an L^∞-measurable $(-1,1)$-form. The holomorphic tangent space is described as the quotient of the complex Banach space of the Beltami differentials (cf. [14, Theorem 7.5]). The holomorphic cotangent space is identified with the space of holomorphic quadratic differentials. The natural pairing between holomorphic tangent space and cotangent space is given by

$$\langle v,q\rangle=\int_M\dot{\mu}q=\int_M\dot{\mu}(z)q(z)\cdot\frac{i}{2}dz\wedge d\overline{z}$$

for $v=[\dot{\mu}]\in T_x\mathcal{T}_{g,m}$ and $q\in T_x^*\mathcal{T}_{g,m}\cong\mathcal{Q}_x$ and $x=(M,f)\in\mathcal{T}_{g,m}$.

Convention 1. *Henceforth, for (p_i,q_i) forms $\varphi_i=\varphi_i(z)dz^{p_i}d\overline{z}^{q_i}$ $(i=1,2)$ on a Riemann surface M with $p_1+p_2=q_1+q_2=1$, we write*

$$\int_M\varphi_1\varphi_2=\int_M\varphi_1(z)\varphi_2(z)\cdot\frac{i}{2}dz\wedge d\overline{z}.$$

$$\begin{tikzcd}
0 & & \\
C^0(\mathcal{U},\Omega_M^{\otimes 2}) \arrow[u] \arrow[r, "\delta"] & C^1(\mathcal{U},\Omega_M^{\otimes 2}) & \\
C^0(\mathcal{U},\Theta_M) \arrow[u, "L.q_0"] \arrow[r, "\delta"] & C^1(\mathcal{U},\Theta_M) \arrow[u, "-L.q_0"] \arrow[r, "\delta"] & C^2(\mathcal{U},\Theta_M)
\end{tikzcd}$$

FIGURE 1. Double complex for the tangent spaces to $\mathcal{Q}_{g,m}$

3. COORDINATES VIA REPRESENTATIONS OF THE ODD COHOMOLOGY

3.1. **The tangent spaces to quadratic differentials.** Following [13], we describe the tangent space $T_{q_0}\mathcal{Q}_{g,m}$ as the first hypercohomolgy group $\mathbb{H}^1(L^\bullet)$ a complex of sheaves (cf. [11] or [12]). We will need the Kodaira-Spencer identification of the tangent space of Teichmüller space with the first cohomology group of the sheaf of holomorphic vector fields (for instance, see [18]. See also [14] and [15]).

Let X and q be a holomorphic vector field and a holomorphic quadratic differential on an open set of a Riemann surface M. Denote by $L_X q$ the Lie derivative of q along X. Let Θ_M and $\Omega_M^{\otimes 2}$ be the sheaves of germs of holomorphic vector fields with zeroes at punctures and meromorphic quadratic differentials on M with (at most) first order poles at punctures, respectively. Let $q_0 \in \mathcal{Q}_{g,m}$ (q_0 need not to be generic). The tangent space $T_{q_0}\mathcal{Q}_{g,m}$ is identified with the first hypercohomology group of the complex of sheaves

$$L^\bullet\colon \quad 0 \longrightarrow \Theta_M \xrightarrow{L.q_0} \Omega_M^{\otimes 2} \longrightarrow 0$$

(cf. [13, Proposition 4.5]). For the convenience, we recall the definition of the (first) hypercohomology group which we use here. The first cochain group is the direct sum $C^0(M,\Omega_M^{\otimes 2}) \oplus C^1(M,\Theta_M)$. Consider an appropriate covering $\mathcal{U}=\{U_i\}_i$ on M such that $\mathbb{H}^1(L^\bullet) \cong \mathbb{H}^1(\mathcal{U},L^\bullet)$ (see the proof of [13, Proposition 4.5]). A cochain $(\{\phi_i\}_i,\{X_{ij}\}_{i,j})$ in $C^0(\mathcal{U},\Omega_M^{\otimes 2}) \oplus C^1(\mathcal{U},\Theta_M)$ is said to be *cocycle* if it satisfies

$$\delta\{X_{ij}\}_{i,j} = X_{ij}+X_{jk}+X_{ki}=0, \quad \delta\{\phi_i\}_i = \phi_i-\phi_j = L_{X_{ij}}(q_0).$$

A *coboundary* is a cochain $(\{\phi_i\}_i,\{X_{ij}\}_{i,j})$ of the form

$$X_{ij} = Z_i - Z_j = \delta\{Z_i\}_i, \quad \phi_i = L_{Z_i}(q_0)$$

for some 0-cochain $\{Z_i\}_i \in C^0(\mathcal{U},\Theta_M)$ (cf. Figure 1).

3.2. **Branched covering space associated with generic differentials.** Let $x=(M,f)\in\mathcal{T}_{g,m}$ and q_0 a generic holomorphic quadratic differential on M. We consider M as a pair of topological surface Σ_g and a complex strtucture on $\Sigma_g - \{\text{punctures of } M\}$.

Let U be a contractible neighborhood of q_0 in $\mathcal{Q}^0_{g,m}$. For $q\in U$, let $A_q = \mathrm{Zero}(q)\cup\mathrm{Pole}(q)\subset\Sigma_g$ and M_q the underlying surface of q. Then, we have a homomorphism

$$\rho_q\colon \pi_1(\Sigma_g - A_q) \to \mathbb{Z}/2\mathbb{Z} \subset \mathrm{Isom}(\mathbb{C}) \tag{3.1}$$

associated to the double covering space

$$\pi_q \colon \tilde{M}_q \to M_q$$

of Riemann surface of $\sqrt{q}$ over $\Sigma_g - A_q$, where $\mathbb{Z}/2\mathbb{Z}$ in (3.1) is recognized as a subgroup of the isometry group $\mathrm{Isom}(\mathbb{C})$ of $\mathbb{C}$ with the standard Euclidean metric generated by the π-rotation. The square root $\sqrt{q}$ lifts to a well-defined holomorphic 1- form ω_q on $\tilde{M}_q$.

Let $W = \Sigma_g - A_{q_0}$ and W' be a compact subsurface of W with smooth boundary such that the inclusion $W' \hookrightarrow W_0$ is homotopy equivalent and the complement $\Sigma_g - W'$ is a union of tiny open $|q_0|$-disks of centers in A_0 By taking sufficiently small U if necessary, we may assume that $A_q \subset \Sigma_g - W'$ for any $q \in U$. Hence, by passing the inclusions $W' \hookrightarrow W$ and $W' \hookrightarrow \Sigma_g - A_q$, we get a canonical identification between $\pi_1(W)$ and $\pi_1(\Sigma_g - A_q)$ so that $\rho_q = \rho_0$ as a homomorphism from $\pi_1(W)$ to $\mathbb{Z}/2\mathbb{Z}$ for all $q \in U$.

3.3. **Backgrounds.** Let $q_0 \in \mathcal{Q}_{g,m}$ be a generic differential on M_{q_0}. Take a neighborhood U of q_0 as the previous section. Then, $\mathbb{M} = \cup_{q \in U} \tilde{M}_q$ is a holomorphic family of closed Riemann surfaces over U, and the arrangement of the covering transformation $r_q \colon \tilde{M}_q \to \tilde{M}_q$ defines the holomorphic automorphism $\mathbf{r}$ of $\mathbb{M}$ of order 2. Taking a sufficiently small U if necessary, we may assume that there is a covering $\{\mathcal{V}_i\}_i$ of $\mathbb{M}$ such that

(1) each $\mathcal{V}_i$ admits a biholomorphic mapping $\mathcal{A}_i$ onto the product domain $\mathbb{D} \times U$ such that its restriction to $\mathcal{V}_{i,q} = \mathcal{V}_i \cap \tilde{M}_q$ is a biholomorphism onto $\mathbb{D} \times \{q\}$ for each $q \in U$,
(2) for any finite $i_1, \cdots, i_k$ and $q \in U$, the intersection $\mathcal{V}_{i_1,q} \cap \cdots \cap \mathcal{V}_{i_k,q}$ is connected and simply connected,
(3) eacn $V_{i,q}$ contains at most one branch point of $\tilde{M}_q \to M_q$ and any branch point of $\tilde{M}_q \to M_q$ is contained in at most one open set in $\{V_{i,q}\}_i$, and
(4) the covering $\{\mathcal{V}_i\}_i$ is equivariant under $\mathbf{r}$ in the sense that $\{\mathbf{r}(\mathcal{V}_i)\}_i = \{\mathcal{V}_i\}_i$ and $\mathcal{A}_j \circ \mathbf{r} = \mathcal{A}_i$ when $\mathbf{r}(\mathcal{V}_i) = \mathcal{V}_j$.

Set $D = \{\lambda = s + it \mid |s|, |t| < \delta_0\}$. Let $\{q_\lambda\}_{\lambda \in D}$ be a differentiable family of holomorphic quadratic differenitals with $q_0 = q_0$ and $q_\lambda \in U$ for $\lambda \in D$. For any $\lambda \in D$, we let $\alpha_i(\lambda) \colon \mathcal{V}_{i,q_\lambda} \to \mathbb{D}$ by $\alpha_i(\lambda) = \mathcal{A}_i \mid_{\tilde{M}_{q_\lambda}}$ and set $V_{\lambda;ij} = \alpha_j(\lambda)(\mathcal{V}_{i,q_\lambda} \cap \mathcal{V}_{j,q_\lambda})$, and $\alpha_{ij}(\lambda) = \alpha_j(\lambda) \circ \alpha_i(\lambda)^{-1} \colon V_{\lambda;ji} \to V_{\lambda;ij}$. For simplicity, set $\alpha_i = \alpha_{0;i}$, $\alpha_{ij} = \alpha_{0;ij}$ and $V_{ij} = V_{0;ij}$. Consider the image of α_j is in the z_j-plane.

Let $\tilde{q}_{\lambda;i} dz_i^2$ be the representation of $(\pi_{q_\lambda})^*(q_\lambda)$ under the coordinate $\alpha_i(\lambda)$. Let

$$\tilde{\phi}_{\lambda;i} = \frac{\partial \tilde{q}_{\lambda;i}}{\partial \lambda}, \quad \tilde{X}_{\lambda;i} = -\frac{\partial \alpha_i(\lambda)}{\partial \lambda} \circ \alpha_i(\lambda)^{-1}, \tag{3.2}$$

on $\mathbb{D}$ for all i and set

$$\tilde{X}_{\lambda;ij}(z_j) = \frac{\partial \alpha_{ij}(\lambda)}{\partial \lambda}(z_i),$$

where $z_j = \alpha_{ij}(\lambda)(z_i)$ and $z_i \in V_{\lambda;ij}$ (tildes mean objects (differentials or vector fields etc.) on the covering space $\tilde{M}_{q_0}$ which obtained as lifts of objects on M_{q_0}). Then,

$$\tilde{X}_{\lambda;i}(z_i)\frac{\partial}{\partial z_i}, \quad \tilde{X}_{\lambda;ij}(z_j)\frac{\partial}{\partial z_j}$$

are vector fields on appropriate sets of $\tilde{M}_{q_\lambda}$. Set $\tilde{q}_i = \tilde{q}_{0;i}$ and $\tilde{\phi}_i = \tilde{\phi}_{0;i}$. Notice that

$$\alpha_{ij}(\lambda)_*(\tilde{X}_{\lambda;i}) - \tilde{X}_{\lambda;j} = \tilde{X}_{\lambda;ij}, \tag{3.3}$$

$$\alpha_{ij}(\lambda)^*(\tilde{\phi}_{\lambda;i}) - \tilde{\phi}_{\lambda;j} = L_{\tilde{X}_{\lambda;ij}}(\tilde{q}_{\lambda;j}) \tag{3.4}$$

on $V_{\lambda;ij}$. We abbreviate these equations to

$$\tilde{X}_{\lambda;i} - \tilde{X}_{\lambda;j} = \tilde{X}_{\lambda;ij}, \quad \tilde{\phi}_{\lambda;i} - \tilde{\phi}_{\lambda;j} = L_{\tilde{X}_{\lambda;ij}}(\tilde{q}_\lambda). \tag{3.5}$$

for instance. Notice again that $\omega_{q_\lambda}^2 = (\pi_{q_\lambda})^*(q_\lambda)$ for all $\lambda \in D$.

Notice that for any cocycle $\{X_{ij}\}_{ij} \in C^1(M, \Theta_M)$, we can always find a 0-cochain $\{X_i\}_i$ of the sheaf of C^∞-vector fields which satisfies $X_i - X_j = X_{ij}$ on $U_i \cap U_j$ applying a partition of unity. Then, from (3.2), the $\overline{z}$-derivative

$$\dot{\nu} = -(\tilde{X}_i)_{\overline{z}} \tag{3.6}$$

defines a Beltrami differential on M which represents the infinitesimal deformation corresponding to the tangent vector associated to the cohomology class of $\{X_{ij}\}_{i,j}$ in $H^1(M, \Theta_M)$. Compare with Equation (7.27) in [14, §7.2.4].

3.4. **Local coordinates via homomorphisms.** Let $H_1(\tilde{M}_q)^-$ is the odd part of the homology of $\tilde{M}_q$ with coefficient in $\mathbb{Z}$. Namely, $H_1(\tilde{M}_q)^-$ is the the eigen space of the eigen value -1 of the linear automorphism $(r_q)_*$ on the first cohomology group of $\tilde{M}_q$ defined by the covering transformation r_q. In [4], Douady and Hubbard consider the following mapping

$$\mathrm{DH}\colon U \ni \lambda \mapsto \chi_q \in \mathrm{Hom}(H_1(\tilde{M}_{q_0})^-, \mathbb{C}) \tag{3.7}$$

defined by

$$\chi_q(c) = \int_c \omega_q$$

where we identify $H_1(\tilde{M}_{q_0})^- \cong H_1(\tilde{M}_q)^-$ in a canonical way as discussed in §3.2. Notice that $\mathrm{Hom}(H_1(\tilde{M}_{q_0})^-, \mathbb{C})$ is a $\mathbb{C}$-vector space of complex dimension $6g-6+2m$. We shall check the following.

Lemma 3.1 (Differential of DH)**.** *Let $q_0 \in \mathcal{Q}_{g,m}$ be a generic differential. Let $v \in T_{q_0}\mathcal{Q}_{g,m}$ be a tangent vector associated to a 1-cochain $(\{\phi_i\}_i, \{X_{ij}\}_{i,j})$. Let $\{X_i\}_i$ be a 0-cochain of the sheaf of C^∞-vector fields on X_{q_0}. Then, the derivative of the mapping* (3.7) *along v satisfies*

$$\mathrm{DH}_*[v](c) = \int_c \Phi \tag{3.8}$$

for all $c \in H_1(\tilde{M}_{q_0})^-$, where

$$\Phi = \left(\frac{\tilde{\phi}_i}{2\omega_{q_0}} - \omega'_{q_0}\tilde{X}_i - \omega_{q_0}(\tilde{X}_i)_z\right)dz - \omega_{q_0}(\tilde{X}_i)_{\overline{z}}d\overline{z}$$

on $\mathcal{V}_{i,0}$.

One can easily check that Φ in Lemma 3.1 is a well-defined closed one-form on $\tilde{M}_{q_0}$ which satisfies $(r_{q_0})^*(\Phi) = -\Phi$.

Proof. We calculate the λ-derivative at $\lambda = 0$ with the notation in §3.3. Notice that

$$\begin{aligned} \alpha_i(\lambda) \circ \alpha_i^{-1}(z_i) &= z_i - \lambda\tilde{X}_i(z_i) - \overline{\lambda}\tilde{Y}_i(z_i) + o(|\lambda|) \\ \tilde{q}_{\lambda;i}(z_i) &= \tilde{q}_i(z_i) + \lambda\tilde{\phi}_i(z_i) + \overline{\lambda}\tilde{\psi}_i(z_i) + o(|\lambda|) \end{aligned}$$

on $\mathbb{D} = \alpha_i(\mathcal{V}_{i,0})$ as $\lambda \to 0$, where

$$\tilde{Y}_i = -\left.\frac{\partial\alpha_i(\lambda)}{\partial\overline{\lambda}}\right|_{\lambda=0} \circ \alpha_i^{-1}, \quad \tilde{\psi}_i = \left.\frac{\partial q_{\lambda;i}}{\partial\overline{\lambda}}\right|_{\lambda=0},$$

and the convergence is valid uniformly on any compact sets of $\mathbb{D}$. In our case, λ-derivative of $D \ni \lambda \mapsto \mathrm{DH}(q_\lambda)(c)$ at $\lambda = 0$ will be the desired formula.

Notice that the partial λ- derivative of ω_λ at $\lambda = 0$ is

$$\left.\frac{\partial\omega_\lambda}{\partial\lambda}\right|_{\lambda=0} = \frac{\tilde{\phi}_i}{2\omega_{q_0}}$$

on $\mathbb{D} = \alpha_i(\mathcal{V}_{i,0})$. Notice also in general that the derivative of a function defined by the integration

$$(-\delta, \delta) \ni s \mapsto \int_{a(s)}^{b(s)} f(s, x)dx$$

at $s = 0$ is equal to

$$\int_{a(0)}^{b(0)} \frac{\partial f}{\partial s}(0, x)dx + f(0, b(0))\frac{db}{ds}(0) - f(0, a(0))\frac{da}{ds}(0).$$

Hence, by the standard argument, like as the discussion in Proposition 1 of [4], one can check the equations (3.8) hold. □

Though the following is well-known (cf. [34], [35], [25]), we confirm for the following for the completeness. Indeed, the calculation in the following proof will be a guide in the later discussion.

Lemma 3.2 (DH defines a local coordinate)**.** *Let* $q_0 \in \mathcal{Q}_{g,m}$ *be as above. When we take the above neighborhood* U *of* q_0 *to be sufficiently small, the mapping* (3.7) *defines a holomorphic local coordinate around* q_0*.*

Proof. Let $v \in T_{q_0}\mathcal{Q}_{g,m}$ be a tangent vector which is represented by the 1 cocycle $(\{\phi_i\}_i, \{X_{ij}\}_{i,j})$ in $C^0(\mathcal{V}_0, \Omega_{M_{q_0}}^{\otimes 2}) \oplus C^1(\mathcal{V}_0, \Theta_{M_{q_0}})$, where $\mathcal{V}_0 = \{\{\pi_{q_0}(\mathcal{V}_{i,0})\}_i\}_i$ is a covering of M_{q_0} (cf. (3.4)).

Suppose that $\mathrm{DH}_*[v] = 0$. Let Φ be a closed one form defined in Lemma 3.1. Since $(r_0)^*(\Phi) = -\Phi$,

$$\int_c \Phi = \int_{r_0(c)} \Phi = -\int_c \Phi$$

for all $c \in H_1(\tilde{M}_{q_0})^+$, where $H_1(\tilde{M}_{q_0})^+$ is the even part of the homology of $\tilde{M}_q$ with coefficient in $\mathbb{Z}$. Hence, there is a smooth function f on $\tilde{M}_{q_0}$ such that $df = \Phi$ from de Rham's theorem. Namely,

$$\begin{aligned} \frac{\tilde{\phi}_i}{2\omega_{q_0}} - \omega'_{q_0}\tilde{X}_i - \omega_{q_0} \cdot (\tilde{X}_i)_z &= f_z \\ -\omega_{q_0} \cdot (\tilde{X}_i)_{\overline{z}} &= f_{\overline{z}} \end{aligned}$$

hold on $\mathbb{D} = \alpha_i(\mathcal{V}_{i,0})$, where $\tilde{\phi}_i$ and $\tilde{X}_i$ are lifts of ϕ_i and $\tilde{X}_i$ to $\tilde{M}_{q_0}$, respectively. Since $\{(\mathcal{V}_{i,\lambda}, \alpha_i(\lambda))\}_i$ is a system of holomorphic local coordinates on $\tilde{M}_{q_\lambda}$ for each $\lambda \in D$, $\dot{\nu} = -(\tilde{X}_i)_{\overline{z}}$ defines a Beltrami differential on $\tilde{M}_{q_0}$ which descends to a Beltrami differential $\dot{\mu}$ on M_{q_0}. Notice that for any holomorphic quadratic differential φ on M_{q_0}, the quotient $(\pi_{q_0})^*(\varphi)/\omega_{q_0}$ is a holomorphic 1-form on $\tilde{M}_{q_0}$ and

$$\begin{aligned}
\int_{M_{q_0}} \dot{\mu}\varphi &= \frac{1}{2}\int_{\tilde{M}_{q_0}} \dot{\nu} \cdot (\pi_{q_0})^*(\varphi) = \frac{1}{2}\int_{\tilde{M}_{q_0}} f_{\overline{z}} \frac{(\pi_{q_0})^*(\varphi)}{\omega_{q_0}} \\
&= \frac{1}{2}\int_{\tilde{M}_{q_0}} \overline{\partial}\left(f(z) \frac{(\pi_{q_0})^*(\varphi)(z)}{\omega_{q_0}(z)} dz \right) \\
&= \frac{1}{2}\int_{\tilde{M}_{q_0}} d\left(f(z) \frac{(\pi_{q_0})^*(\varphi)(z)}{\omega_{q_0}(z)} dz \right) = 0.
\end{aligned}$$

Hence, $\dot{\mu}$ is an infinitesimally trivial Beltrami differenital on M_{q_0} by Teichmüller lemma (cf. Lemma 7.6 of [14]). This means that the cohomology class

$$[\{X_{ij}\}_{i,j}] \in H^1(\mathcal{V}_0, \Theta_{M_{q_0}}) \cong H^1(M_{q_0}, \Theta_{M_{q_0}})$$

is trivial from (3.5) since $\mathcal{V}_0$ is a Leray covering (see also Theorem 3.5 of [18]). Therefore, there is a 0-cochain $\{Z_i\}_i \in C^0(\mathcal{V}_{0,i}, \Theta_{M_{q_0}})$ such that $Z_i - Z_j = X_{ij}$ for all i, j.

Let $\zeta_i = L_{Z_i}(q_0) \in \Gamma(\mathcal{V}_{0,i}, \Omega_{M_{q_0}}^{\otimes 2})$ and $\tilde{\zeta}_i = L_{\tilde{Z}_i}(\omega_{q_0}^2)$ be the lift of ζ_i. Then,

$$\tilde{\zeta}_i - \tilde{\zeta}_j = L_{\tilde{X}_{ij}}(\omega_{q_0}^2)$$

and hence $\{\tilde{\phi}_i - \tilde{\zeta}_i\}_i$ defines a holomorphic quadratic differential $\tilde{\varphi}$ on $\tilde{M}_{q_0}$ which descends to a holomorphic quadratic differential φ on M_{q_0} associated to $\{\phi_i - \zeta_i\}_i$. Since $\tilde{X}_i - \tilde{Z}_i = \tilde{X}_j - \tilde{Z}_j$ on $\mathcal{V}_{i,0} \cap \mathcal{V}_{j,0}$, $\{\tilde{X}_i - \tilde{Z}_i\}_i$ defines a vector field $\tilde{W}$ on $\tilde{M}_{q_0}$ which satisfies

$$\begin{aligned}
&((\omega_{q_0})'(\tilde{X}_i) + \omega_{q_0} \cdot (\tilde{X}_i)_z)dz + \omega_{q_0} \cdot (\tilde{X}_i)_{\overline{z}} d\overline{z} - \frac{L_{\tilde{Z}_i}(\omega_{q_0}^2)}{2\omega_{q_0}} \\
&= \left((\omega_{q_0})'(\tilde{X}_i - \tilde{Z}_i) + \omega_{q_0} \cdot (\tilde{X}_i - \tilde{Z}_i)_z\right) dz + \omega_{q_0} \cdot (\tilde{X}_i - \tilde{Z}_i)_{\overline{z}} d\overline{z} \\
&= d(\omega_{q_0} \cdot \tilde{W})
\end{aligned}$$

since each $\tilde{Z}_i$ is a holomorphic section. Consequently, we deduce from the assumption that

$$0 = \int_c \Omega = \int_c \left(\Omega - \frac{\tilde{\zeta}_i - L_{\tilde{Z}_i}(\omega_{q_0}^2)}{2\omega_{q_0}}\right) = \int_c \frac{\tilde{\varphi}}{2\omega_{q_0}} - \int_c d(\omega_{q_0} \cdot W) = \int_c \frac{\tilde{\varphi}}{2\omega_{q_0}} \tag{3.9}$$

for all $c \in H_1(\tilde{M}_{q_0})$, and hence $\tilde{\varphi} = 0$ and $\varphi = 0$. Therefore $\phi_i = \zeta_i$ and

$$(\{\phi_i\}_i, \{X_{ij}\}_{i,j}) = (\{L_{Z_i}(q_0)\}_i, \{Z_i - Z_j\}_{i,j}).$$

Thus, the cohomology class of the cochain $(\{\phi_i\}_i, \{X_{ij}\}_{i,j})$ vanishes.

As a consequence, the derivative of the map DH on the tangent space $T_{q_0}\mathcal{Q}_{g,m} \cong \mathbb{H}^1(L^\bullet)$ is injective. Since the dimensions of $\mathcal{Q}_{g,m}$ and $\mathrm{Hom}(H_1(\tilde{M}_{q_0})^-, \mathbb{C})$ are same, the derivative of DH is an isomorphism from $T_{q_0}\mathcal{Q}_{g,m}$ onto $\mathrm{Hom}(H_1(\tilde{M}_{q_0})^-, \mathbb{C})$. □

3.5. **Homomorphisms on slices.** For $F \in \mathcal{MF} - \{0\}$. we define a slice

$$E^v(F) = \{q \in \mathcal{Q}_{g,m} \mid \mathbf{v}(q) = F\}.$$

for F in $\mathcal{Q}_{g,m}$. Hubbard and Masur showed that the restriction of the projection

$$E^v(F) \ni q \mapsto \pi(q) \in \mathcal{T}_{g,m}$$

is a homeomorphism ([13, §2]). The following two lemmas are well-known (cf. [13, Lemma 4.3] and [24, Proposition 1]. See also [6]).

Lemma 3.3. *Let $q_0 \in \mathcal{Q}_{g,m}$ be a generic differential. When we take U sufficiently small,*

$$E^v(\mathbf{v}(q_0)) \cap U = \{q \in \mathcal{Q}_{g,m} \mid \mathrm{Re}(\mathrm{DH}(q)) = \mathrm{Re}(\mathrm{DH}(q_0))\}$$

holds.

Lemma 3.4. *Let $x_0 \in \mathcal{T}_{g,m}$ and $F \in \mathcal{MF}$. If q_{F,x_0} is generic, then the map*

$$\mathcal{T}_{g,m} \ni x \mapsto q_{F,x} \in \mathcal{Q}_{g,m}$$

is real analytic around x_0.

4. Family of quadratic differentials with prescribed vertical foliation

Fix $F \in \mathcal{MF}$. Consider a family $\{q_\lambda\}_{\lambda \in D}$ of holomorphic quadratic differentials such that $\mathbf{v}(q_\lambda) = F$. Suppose that q_0 is generic and $x_\lambda = \pi(q_\lambda) \in \mathcal{T}_{g,m}$ varies holomorphically. Then, the family $\{q_\lambda\}_{\lambda \in D}$ is a real analytic family of quadratic differentials (cf. Lemma 3.4). Take a neighborhood U of q_0 as the previous sections.

4.1. **Differential via the complex conjugate.** We first notice a simple observation. Let $\pi\colon E \to M$ be a holomorphic vector bundle. Let $g\colon \mathbb{D} \to M$ be a holomorphic mapping and $G\colon \mathbb{D} \to E$ be a C^1 mapping such that $\pi \circ G = g$ on $\mathbb{D}$. Then,

$$\pi_*\left(G_*\left(\frac{\partial}{\partial \overline{\lambda}}\right)\right) = g_*\left(\frac{\partial}{\partial \overline{\lambda}}\right) = 0$$

since g is holomorphic. Hence,

$$G_*\left(\frac{\partial}{\partial \overline{\lambda}}\right)\bigg|_{\lambda} \in \mathrm{Ker}(\pi_*) \cong E_{g(\lambda)} \tag{4.1}$$

for all $\lambda \in \mathbb{D}$.

Lemma 4.1 (First derivatives)**.** *Let $\{q_\lambda\}_{\lambda \in D}$ be the family of holomorphic quadratic differentials defined as above. Let η_λ be a holomorphic quadratic differential on X_{q_λ} such that the infinitesimal Bertrami differential $\dot{\mu}$ associated to the tangent vector for the λ-derivative of the mapping*

$$D \ni \lambda \mapsto \pi(q_\lambda) \in \mathcal{T}_{g,m}$$

at λ satisfies

$$\int_{M_{q_\lambda}} \dot{\mu}\psi = \int_{M_{q_\lambda}} \frac{\overline{\eta_\lambda}}{|q_\lambda|}\psi$$

for all holomorphic quadratic differential ψ on M_{q_λ}. Then,

$$\frac{\partial\, \chi_{q_\lambda}(c)}{\partial \lambda} = \int_c \overline{\left(\frac{(\pi_{q_\lambda})^*(\eta_\lambda)}{\omega_{q_\lambda}}\right)}, \quad \frac{\partial\, \chi_{q_\lambda}(c)}{\partial \overline{\lambda}} = -\int_c \frac{(\pi_{q_\lambda})^*(\eta_\lambda)}{\omega_{q_\lambda}}$$

for any $c \in H_1(\tilde{M}_{q_0})$ and $\lambda \in D$.

Proof. The existence and the uniqueness of η_λ follows from an argumant by David Dumas [6, Theorem 5.3] since each q_λ is generic. Indeed, the pairing

$$(\eta, \psi) \mapsto \int_{M_{q_\lambda}} \frac{\overline{\eta}\psi}{|q_\lambda|}$$

is a positive definite hermitian inner product on the space of holomorphic quadratic differentials on M_{q_λ}. Notice that

$$\frac{\partial\, \chi_{q_\lambda}(c)}{\partial\overline{\lambda}} = 0 = \int_c \frac{(\pi_{q_\lambda})^*(\eta_\lambda)}{\omega_{q_\lambda}}$$

for any $c \in H_1(\tilde{M}_{q_0})^+$ and $\lambda \in D$. It suffices to show the case where $\lambda = 0$ and $c \in H_1(\tilde{M}_{q_0})^-$. As we discussed around (4.1), from Lemma 3.1 and the proof of Lemma 3.2, there is a holomorphic quadratic differential φ on X_{q_0} such that

$$\left.\frac{\partial\, \chi_{q_\lambda}(c)}{\partial\overline{\lambda}}\right|_{\lambda=0} = \int_c \frac{(\pi_{q_0})^*(\varphi)}{\omega_{q_0}}$$

for any $c \in H_1(\tilde{M}_{q_0})^-$ (see (3.9)). Since the vertical foliation $|\mathrm{Re}(\omega_{q_\lambda})|$ of ω_{q_λ} is unchanged on D, for all $c \in H_1(\tilde{M}_{q_0})^-$, the real part of $\chi_{q_\lambda}(c)$ is a constant function on D by Lemma 3.3. Therefore,

$$\frac{\partial\, \chi_{q_\lambda}(c)}{\partial\overline{\lambda}} = -\overline{\frac{\partial\, \chi_{q_\lambda}(c)}{\partial\lambda}}$$

on D. Let $(\{\phi_i\}_i, \{X_{ij}\}_{i,j})$ be the cocycle representing the tangent vector associated to the λ-derivative of the mapping $D \ni \lambda \mapsto q_\lambda$ at $\lambda = 0$. Let $\{X_i\}_i$ be a 0-cochain of the sheaf of C^∞-vector fields with $X_i - X_j = X_{ij}$. Then,

$$\Omega = \left(\frac{\tilde{\phi}_i}{2\omega_{q_0}} - \omega'_{q_0}\tilde{X}_i - \omega_{q_0}(\tilde{X}_i)_z\right)dz - (\omega_{q_0}(\tilde{X}_i)_{\overline{z}})d\overline{z}$$

on U_i defines a closed 1-form on $\tilde{M}_{q_0}$ satisfies

$$\frac{\partial\, \chi_{q_\lambda}(c)}{\partial\lambda} = \int_c \Omega$$

for all $c \in H_1(\tilde{M}_{q_0})$. Therefore, there is a C^∞-function f on $\tilde{M}_{q_0}$ such that

$$\Omega = -\overline{\frac{(\pi_{q_0})^*(\varphi)(z)}{\omega_{q_0}(z)}}d\overline{z} + df,$$

that is,

$$\frac{\tilde{\phi}_i}{2\omega_{q_0}} - \omega'_{q_0}\tilde{X}_i - \omega_{q_0}(\tilde{X}_i)_z = f_z \tag{4.2}$$

$$-\omega_{q_0}(\tilde{X}_i)_{\overline{z}} = -\overline{\left(\frac{(\pi_{q_0})^*(\varphi)(z)}{\omega_{q_0}(z)}\right)} + f_{\overline{z}} \tag{4.3}$$

on $\mathcal{V}_{i,0}$. From (4.3), the lift $(\pi_{q_0})^*(\dot{\mu})$ of $\dot{\mu} = -(X_i)_{\overline{z}}$ satisfies

$$(\pi_{q_0})^*(\dot{\mu}) = -\frac{\overline{(\pi_{q_0})^*(\varphi)}}{|\omega_{q_0}|^2} + \frac{f_{\overline{z}}}{\omega_{q_0}} \tag{4.4}$$

on $\tilde{M}_{q_0}$. Let ψ be a holomorphic quadratic differenital on M_{q_0}. From (4.4)

$$\begin{aligned}\int_{\tilde{M}_{q_0}}(\pi_{q_0})^*(\dot{\mu})(\pi_{q_0})^*(\psi) &= -\int_{\tilde{M}_{q_0}}\frac{\overline{(\pi_{q_0})^*(\varphi)}}{|\omega_{q_0}|^2}(\pi_{q_0})^*(\psi)+\int_{\tilde{M}_{q_0}}\frac{f_{\overline{z}}}{\omega_{q_0}}(\pi_{q_0})^*(\psi)\\ &= -\int_{\tilde{M}_{q_0}}\frac{\overline{(\pi_{q_0})^*(\varphi)}}{|\omega_{q_0}|^2}(\pi_{q_0})^*(\psi)+\int_{\tilde{M}_{q_0}}d\left(f\frac{(\pi_{q_0})^*(\psi)}{\omega_{q_0}}\right)\\ &= -\int_{\tilde{M}_{q_0}}\frac{\overline{(\pi_{q_0})^*(\varphi)}}{|\omega_{q_0}|^2}(\pi_{q_0})^*(\psi)\end{aligned}$$

By descending to M_{q_0}, we have

$$\int_{M_{q_0}}\frac{\overline{\eta_0}}{|q_0|}\psi=\int_{M_{q_0}}\dot{\mu}\psi=-\int_{M_{q_0}}\frac{\overline{\varphi}\psi}{|q_0|}$$

for all ψ. Therefore, $\varphi=-\eta_0$ from [6, Theorem 5.3] again. □

4.2. **Laplacian of homomorphisms.** This section is devoted to calculating the Laplacian of Douady-Hubbard map (3.7). The author must confess that for the proof of the plurisubharmonicity of extremal length funcitons, we need the equation (4.6) rather than the formula (4.5) in the following lemma. However, we give the following lemma for its own interests.

Lemma 4.2 (Laplacian)**.** *Under the assumption in Lemma 4.2, we have*

$$\left.\frac{\partial^2(\chi_{q_\lambda}(c))}{\partial\lambda\partial\overline{\lambda}}\right|_{\lambda=0}=-4i\,\mathrm{Im}\int_{\tilde{M}_{q_0}}\frac{|(\pi_{q_0})^*(\eta_0)|^2}{|\omega_{q_0}|^2}\frac{\sigma_c}{\omega_{q_0}}\tag{4.5}$$

for $c\in H_1(\tilde{M}_{q_0})$, where $\sigma_c=\sigma_c(z)dz$ is the holomorphic part of the reproducing harmonic differential associated to c.

Proof. When the homology class c is in the even homology group, $\chi_{q_\lambda}(c)=0$. The integrand of the right-hand side of (4.5) satisfies

$$r_0^*\left(\frac{|(\pi_{q_0})^*(\eta_0)|^2}{|\omega_{q_0}|^2}\frac{\sigma_c}{\omega_{q_0}}\right)=\frac{|(\pi_{q_0})^*(\eta_0)|^2}{|-\omega_{q_0}|^2}\frac{\sigma_c}{-\omega_{q_0}}=-\frac{|(\pi_{q_0})^*(\eta_0)|^2}{|\omega_{q_0}|^2}\frac{\sigma_c}{\omega_{q_0}}$$

and hence, the integral over $\tilde{M}_{q_0}$ vanishes. This means that (4.5) holds for homology classes in the even homology group.

By applying the same argument as that of Lemma 3.1, one obtain

$$\left.\frac{\partial^2\,\chi_{q_\lambda}(c)}{\partial\lambda\partial\overline{\lambda}}\right|_{\lambda=0}=\int_c\Omega_0$$

for any $c\in H_1(\tilde{M}_{q_0})$, where Ω_0 is a closed differential of the form

$$\left(\dot{\eta}_0+\left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)'(\tilde{X}_i)+\left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)(\tilde{X}_i)_z\right)dz+\left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)(\tilde{X}_i)_{\overline{z}}d\overline{z}$$

and

$$\dot{\eta}_0=-\left.\frac{\partial}{\partial\lambda}\left(\frac{(\pi_{q_\lambda})^*(\eta_\lambda)}{\omega_{q_\lambda}}\right)\right|_{\lambda=0}.$$

Since the real part of the function $D \ni \lambda \mapsto \chi_{q_\lambda}(c)$ is constant and the Laplacian is the real operator,

$$\int_c \Omega_0 = \left.\frac{\partial^2 \chi_{q_\lambda}(c)}{\partial\lambda\partial\overline{\lambda}}\right|_{\lambda=0} = -\overline{\left.\frac{\partial^2 \chi_{q_\lambda}(c)}{\partial\lambda\partial\overline{\lambda}}\right|_{\lambda=0}} = -\overline{\left(\int_c \Omega_0\right)} = -\int_c \overline{\Omega_0}$$

for all $c \in H_1(\tilde{M}_{q_0})$. Hence, there is a C^∞-function f on $\tilde{M}_{q_0}$ such that

$$\Omega_0 = -\overline{\Omega_0} + df.$$

Namely, we have

$$\begin{aligned}\dot{\eta}_0 + \left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)'(X_i) + \left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)(\tilde{X}_i)_z &= -\overline{\left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)(\tilde{X}_i)_{\overline{z}}} + f_z \\ &= \overline{\left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)}(\pi_{q_0})^*(\dot{\mu}) + f_z,\end{aligned}$$

where $\dot{\mu} = -(X_i)_{\overline{z}}$ is the infinitesimal Beltrami differential representing the tangent vector of the mapping $D \ni\mapsto \pi(q_\lambda)$ at $\lambda = 0$ (cf. (3.6)). Applying the above equation, we also obtain

$$\begin{aligned}\Omega_0 &= \left(\overline{\left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)(\pi_{q_0})^*(\dot{\mu})} + f_z\right)dz - \left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)(\pi_{q_0})^*(\dot{\mu})d\overline{z} \\ &= \overline{\left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)(\pi_{q_0})^*(\dot{\mu})}dz - \left(\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\right)(\pi_{q_0})^*(\dot{\mu})d\overline{z} \\ &\qquad + \frac{1}{2}df + \frac{i}{2}{}^*df. \end{aligned} \tag{4.6}$$

Let $\sigma(c) = \sigma_c(z)dz + \overline{\sigma_c(z)}d\overline{z}$ be the reproducing harmonic differential on $\tilde{M}_{q_0}$ associated to $c \in H_1(\tilde{M}_{q_0})$, where $\sigma_c(z)dz$ is a holomorphic 1-form on $\tilde{M}_{q_0}$ (we call this holomorphic 1-form the *holomorphic part* of $\sigma(c)$). Namely, the differential $\sigma(c)$ is a unique harmonic differential on $\tilde{M}_{q_0}$ which satisfies

$$\int_c \omega = \int_{\tilde{M}_{q_0}} \omega \wedge {}^*\overline{\sigma(c)}$$

for all closed 1-form ω on $\tilde{M}_{q_0}$. Then, by applying the orthogonal decomposition theorem for the space of L^2-closed forms, we have

$$\begin{aligned}\left.\frac{\partial^2 \chi_{q_\lambda}(c)}{\partial\lambda\partial\overline{\lambda}}\right|_{\lambda=0} &= \int_c \Omega_0 = \int_{\tilde{M}_{q_0}} \Omega_0 \wedge {}^*\overline{\sigma(c)} \\ &= \int_{\tilde{M}_{q_0}} \left(\overline{\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}}(\pi_{q_0})^*(\dot{\mu})\,dz - \frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}(\pi_{q_0})^*(\dot{\mu})\,d\overline{z}\right) \\ &\qquad\qquad \wedge {}^*\overline{(\sigma_c(z)dz + \overline{\sigma_c(z)}d\overline{z})} \\ &= \int_{\tilde{M}_{q_0}} \left(\overline{\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}}(\pi_{q_0})^*(\dot{\mu})\,dz - \frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}(\pi_{q_0})^*(\dot{\mu})\,d\overline{z}\right) \\ &\qquad\qquad \wedge (-i\overline{\sigma_c}d\overline{z} + i\sigma_c dz) \\ &= -4i\,\mathrm{Im}\int_{\tilde{M}_{q_0}} \frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}(\pi_{q_0})^*(\dot{\mu})\sigma_c.\end{aligned}$$

Now, we suppose that the homology class c is in $H_1(\tilde{M}_{q_0})^-$. Since $(r_0)_*(c) = -c$, $r_0^*\sigma(c) = -\sigma(c)$. This means that $\sigma(c)$ has zeroes at each branch points of $\pi_{q_0}\colon \tilde{M}_{q_0} \to M_{q_0}$ and the holomorphic quadratic differential

$$\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\sigma_c = \frac{(\pi_{q_0})^*(\eta_0)(z)}{\omega_{q_0}(z)}\sigma_c(z)dz^2$$

descends to a holomorphic quadratic differential on M_{q_0}. Therefore, we obtain from the definition of η_0 that

$$\begin{aligned}\left.\frac{\partial^2\,\chi_{q_\lambda}(c)}{\partial\lambda\partial\overline{\lambda}}\right|_{\lambda=0} &= -4i\,\mathrm{Im}\int_{\tilde{M}_{q_0}}\frac{(\pi_{q_0})^*(\eta_0)}{\omega_{q_0}}\frac{\overline{(\pi_{q_0})^*(\eta_0)}}{|\omega_{q_0}|^2}\sigma_c \\ &= -4i\,\mathrm{Im}\int_{\tilde{M}_{q_0}}\frac{|(\pi_{q_0})^*(\eta_0)|^2}{|\omega_{q_0}|^2}\frac{\sigma_c}{\omega_{q_0}} \qquad (4.7)\end{aligned}$$

(cf. Lemma 4.1). This is what we wanted. □

5. Levi forms and Plurisubharmonicity of subspecies

5.1. **Levi forms of extremal length functions.** For a real-valued C^2-function U on a complex manifold N, the *Levi form* of U is an hermitian inner product on the holomorphic tangent bundle TN of N defined as

$$\mathcal{L}(U)[v,\overline{v}] = \sum_{i,j=1}^{n}\frac{\partial^2 U}{\partial z_i\partial\overline{z}_j}v_i\overline{v_j}$$

for $v = \sum_{i=1}^{n} v_i(\partial/\partial z_i) \in TN$. Let $p \in N$ and $v \in T_pN$. Let $F\colon \mathbb{D} \to N$ be a holomorphic mapping with $f(0) = p$ and $f_*(\partial/\partial\lambda\mid_{\lambda=0}) = v$, then

$$\mathcal{L}(U)[v,\overline{v}] = \left.\frac{\partial^2(U\circ f)}{\partial\lambda\partial\overline{\lambda}}\right|_{\lambda=0}.$$

Theorem 5.1 (Levi forms)**.** *Let $x_0 = (M_0, f_0) \in \mathcal{T}_{g,m}$ and $v = [\dot{\mu}] \in T_{x_0}\mathcal{T}_{g,m}$. Let $F \in \mathcal{MF}$. Suppose that the Hubbard-Masur differential $q_0 = q_{F,x_0}$ is generic. Then, the extremal length function is real analytic around x_0, and the Levi form of the extremal length function $\mathcal{T}_{g,m} \ni x \mapsto \mathrm{Ext}_x(F)$ at x_0 satisfies*

$$\mathcal{L}(\mathrm{Ext}.(F))[v,\overline{v}] = 2\int_{M_0}\frac{|\eta_v|^2}{|q_0|},$$

where η_v is a unique holomorphic quadratic differential on $M_0 = M_{q_0}$ satisfying that

$$\langle v,\psi\rangle = \int_{M_0}\dot{\mu}\psi = \int_{M_0}\frac{\overline{\eta_v}}{|q_0|}\psi$$

for all $\psi \in \mathcal{Q}_{x_0}$.

Remark 5.1. *In §6, we will discuss a geometric interpretation of the anti-complex linear map*

$$T_{x_0}\mathcal{T}_{g,m} \ni v \mapsto \eta_v \in \mathcal{Q}_{x_0}.$$

Proof of Theorem 5.1. We continue to use the symbols defined in the previous sections. Let $\tilde{g} = 4g - 2 + m$ (≥ 2) be the genus of $\tilde{M}_{q_0}$. Let $\{\alpha_k, \beta_k\}_{k=1}^{\tilde{g}}$ be a symplectic generator of $H_1(\tilde{M}_{q_0})$. Let $x_\lambda = \pi(q_\lambda)$ for instance. By the bilinear relation, we have

$$\begin{aligned}\mathrm{Ext}_{x_\lambda}(F) &= \frac{1}{2}\|\omega_{q_\lambda}\|^2 = \frac{1}{4}\int_{\tilde{M}_{q_\lambda}} \omega_{q_\lambda} \wedge {}^*\overline{\omega_{q_\lambda}} \\ &= \frac{i}{4}\sum_{k=1}^{\tilde{g}}\left(\chi_{q_\lambda}(\alpha_k)\overline{\chi_{q_\lambda}(\beta_k)} - \chi_{q_\lambda}(\beta_k)\overline{\chi_{q_\lambda}(\alpha_k)}\right)\end{aligned}$$

for $\lambda \in D$ (cf. [8, Corollary in §III.2.3]). From Lemmas 3.2 and 3.3, $\chi_{q_{F,x}}(\alpha_k)$ and $\chi_{q_{F,x}}(\beta_k)$ are real analytic around x_0 for each k, and hence so is the extremal length function. From Lemma 4.2, we have

$$\begin{aligned}&\frac{\partial^2}{\partial\lambda\partial\overline{\lambda}}\left(\chi_{q_\lambda}(c_1)\overline{\chi_{q_\lambda}(c_2)}\right)\Big|_{\lambda=0} \\ &= \frac{\partial^2\chi_{q_\lambda}(c_1)}{\partial\lambda\partial\overline{\lambda}}\Big|_{\lambda=0}\overline{\chi_{q_0}(c_2)} + \chi_{q_0}(c_1)\,\frac{\partial^2\overline{\chi_{q_\lambda}(c_2)}}{\partial\lambda\partial\overline{\lambda}}\Big|_{\lambda=0} \\ &\quad + \frac{\partial\chi_{q_\lambda}(c_1)}{\partial\lambda}\Big|_{\lambda=0}\frac{\partial\overline{\chi_{q_\lambda}(c_2)}}{\partial\overline{\lambda}}\Big|_{\lambda=0} + \frac{\partial\chi_{q_\lambda}(c_1)}{\partial\overline{\lambda}}\Big|_{\lambda=0}\frac{\partial\overline{\chi_{q_\lambda}(c_2)}}{\partial\lambda}\Big|_{\lambda=0} \\ &= \frac{\partial^2\chi_{q_\lambda}(c_1)}{\partial\lambda\partial\overline{\lambda}}\Big|_{\lambda=0}\overline{\chi_{q_0}(c_2)} + \chi_{q_0}(c_1)\,\frac{\partial^2\overline{\chi_{q_\lambda}(c_2)}}{\partial\lambda\partial\overline{\lambda}}\Big|_{\lambda=0} \\ &\quad + \frac{\partial\chi_{q_\lambda}(c_1)}{\partial\lambda}\Big|_{\lambda=0}\overline{\frac{\partial\chi_{q_\lambda(c_2)}}{\partial\lambda}\Big|_{\lambda=0}} + \overline{\frac{\partial\chi_{q_\lambda}(c_1)}{\partial\lambda}\Big|_{\lambda=0}}\frac{\partial\chi_{q_\lambda}(c_2)}{\partial\lambda}\Big|_{\lambda=0}\end{aligned}$$

for $c_1, c_2 \in H_1(\tilde{M}_{q_0})$, since the real part of $\chi_{q_\lambda}(c)$ is constant for all $c \in H_1(\tilde{M}_{q_0})$ by Lemma 3.3. Notice that the sum of the last two terms of above calculation is a real number. Therefore, we obtain

$$\begin{aligned}&\frac{\partial^2}{\partial\lambda\partial\overline{\lambda}}\left(\chi_{q_\lambda}(\alpha_k)\overline{\chi_{q_\lambda}(\beta_k)} - \chi_{q_\lambda}(\beta_k)\overline{\chi_{q_\lambda}(\alpha_k)}\right)\Big|_{\lambda=0} \\ &= \int_{\alpha_k}\Omega_0\overline{\int_{\beta_k}\omega_{q_0}} + \overline{\int_{\beta_k}\Omega_0}\int_{\alpha_k}\omega_{q_0} - \int_{\beta_k}\Omega_0\overline{\int_{\alpha_k}\omega_{q_0}} - \int_{\alpha_k}\overline{\Omega_0}\int_{\beta_k}\omega_{q_0} \\ &= \int_{\alpha_k}\Omega_0\int_{\beta_k}\overline{\omega_{q_0}} - \int_{\beta_k}\Omega_0\int_{\alpha_k}\overline{\omega_{q_0}} + \int_{\beta_k}\overline{\Omega_0}\int_{\alpha_k}\omega_{q_0} - \int_{\alpha_k}\overline{\Omega_0}\int_{\beta_k}\omega_{q_0}\end{aligned}$$

where Ω_0 is a closed form defined as (4.6). Since ω_{q_0} is a holomorphic differential, ${}^*\omega_{q_0}$ and ${}^*\overline{\omega_{q_0}}$ are harmonic differentials and they are perpendicular to the exact and co-exact differentials in the L^2-inner product of differential forms (cf. [8, §II.3]).

From (4.6), we obtain

$$\begin{aligned}\mathcal{L}(\mathrm{Ext.}(F))[v,\overline{v}] &= \frac{\partial^2}{\partial\lambda\partial\overline{\lambda}}\mathrm{Ext}_{x_\lambda}(F)\Big|_{\lambda=0} \\ &= \frac{i}{4}\sum_{k=1}^{\tilde{g}}\left(\int_{\alpha_k}\Omega_0\int_{\beta_k}\overline{\omega_{q_0}} - \int_{\beta_k}\Omega_0\int_{\alpha_k}\overline{\omega_{q_0}}\right) \\ &\quad + \frac{i}{4}\sum_{k=1}^{\tilde{g}}\left(\int_{\beta_k}\overline{\Omega_0}\int_{\alpha_k}\omega_{q_0} - \int_{\alpha_k}\overline{\Omega_0}\int_{\beta_k}\omega_{q_0}\right) \\ &= \frac{i}{4}\int_{\tilde{M}_{q_0}}\Omega_0\wedge\overline{\omega_{q_0}} - \frac{i}{4}\int_{\tilde{M}_{q_0}}\overline{\Omega_0}\wedge\omega_{q_0} \\ &= \mathrm{Re}\int_{\tilde{M}_{q_0}}\frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}}(\pi_{q_0})^*(\dot{\mu})\omega_{q_0} \\ &= 2\mathrm{Re}\int_{M_{q_0}}\eta_v\dot{\mu} = 2\int_{M_0}\frac{|\eta_v|^2}{|q_0|}\end{aligned}$$

from the definition of the differential η_v (cf. [8, Proposition III.2.3]). This implies what we wanted. □

From Theorem 5.1, we deduce the following inequality.

Corollary 5.1 (Gradients and Levi forms)**.** *Let* $x_0 = (M_0, f_0) \in \mathcal{T}_{g,m}$ *and* $v = [\dot{\mu}] \in T_{x_0}\mathcal{T}_{g,m}$. *Let* $F \in \mathcal{MF}$. *Suppose that* $q_0 = q_{F,x_0}$ *is generic. Then,*

$$2\left|(\mathrm{Ext.}(F))_*[v]\right|^2 \leq \mathrm{Ext}_{x_0}(F)\cdot\mathcal{L}(\mathrm{Ext.}(F))[v,\overline{v}]$$

holds.

Proof. It follows from Gardiner's formula that

$$(\mathrm{Ext.}(F))_*[v] = -\int_{M_0}\dot{\mu}\,q_0 = -\int_{M_0}\frac{\overline{\eta_v}}{|q_0|}q_0. \tag{5.1}$$

See [9, §11.8] (See also §7.1). Notice in comparing the Gardiner's original formula with (5.1) that our differential $q_0 = q_{F,x_0}$ has F as the vertical foliation, while Gardiner considers the quadratic differential with horizontal foliation F. Hence, we have to put the minus sign in (5.1). From Cauchy-Schwarz inequality, one can see from Theorem 5.1 that

$$\begin{aligned}&\mathrm{Ext}_{x_0}(F)\mathcal{L}(\mathrm{Ext.}(F))[v,\overline{v}] - 2\left|(\mathrm{Ext.}(F))_*[v]\right|^2 \\ &= \mathrm{Ext}_{x_0}(F)\cdot 2\int_{M_0}\frac{|\eta_v|^2}{|q_0|} - 2\left|-\int_{M_0}\frac{\overline{\eta_v}}{|q_0|}q_0\right|^2 \\ &= 2\left(\int_{M_0}|q_0|\cdot\int_{M_0}\frac{|\eta_v|^2}{|q_0|} - \left|\int_{M_0}\frac{\overline{\eta_v}}{|q_0|^{1/2}}\frac{q_0}{|q_0|^{1/2}}\right|^2\right) \geq 0,\end{aligned}$$

which implies what we wanted. □

5.2. **Plurisubharmonicity for subspecies.** Let N be a complex manifold. A function U on N is said to be *plurisubharmonic* if U is upper semi-continuous, and $U\circ f$ is subharmonic for all holomorphic mapping $f\colon \mathbb{D}\to N$. A C^2-function on N is plurisubharmonic if the Levi form is positive semi-definite. We call a C^2-function *strictly plurisubharmonic* if the Levi form is positive-definite. When a function U

is plurisubharmonic, so is $g \circ U$ for any increasing convex function g on $\mathbb{R}$ such that the limit $\lim_{t\to-\infty} g(t)$ exists.

In the proof of the following theorem, we remind that if $U(z)$ is a real-valued positive C^2-function on a domain on $\mathbb{C}$, $\log U$ and $-1/U$ satisfy

$$(\log U)_{\lambda\overline{\lambda}} = \frac{U \cdot U_{\lambda\overline{\lambda}} - |U_\lambda|^2}{U^2} \tag{5.2}$$

$$\left(-\frac{1}{U}\right)_{\lambda\overline{\lambda}} = \frac{U \cdot U_{\lambda\overline{\lambda}} - 2|U_\lambda|^2}{U^3}. \tag{5.3}$$

Especially, for a positive C^2-function U on a complex manifold, if $-1/U$ is plurisubharminic, $\log U$ and U are also plurisubharmonic. From (5.2) and (5.3), Theorem 1.1 follows from the following theorem.

Theorem 5.2 (Plurisuperharmonicity of reciprocals). *Let $F_1, F_2, \cdots, F_n$ be measured foliations. Let $a_k, c \geq 0$ $(k = 1, \cdots, n)$ with $\sum_{k=1}^{n} a_k > 0$. If $n \geq 2$, $a_k, c > 0$ $(k = 1, \cdots, n)$ and some two of them are transverse in the sense of* (1.1), *the function*

$$\rho(x) = -\frac{1}{c + \sum_{k=1}^{n} a_k \mathrm{Ext}_x(F_k)}$$

is a negative, plurisubharmonic exhaustion function of $\mathcal{T}_{g,m}$ with lower bound. When $n \geq 1$ or $c \geq 0$, ρ is a negative, plurisubharmonic function on $\mathcal{T}_{g,m}$ with lower bound.

Proof. We only show the case $n = 2$. The other case can be treated in the same way. Furthermore, we may also assume that each F_k are essentially complete, and F_1 and F_2 are transverse in the sense of (1.1). The general case follows from a standard approximating procedure (cf. [17, Theorem 2.6.1], [21, Theorem 2 in Chapter II] or [30, 3.3 in Chapter II]). Under the assumption, each extremal length function $\mathcal{T}_{g,m} \ni x \mapsto \mathrm{Ext}_x(F_k)$ is real analytic.

Let $x = (M, f) \in \mathcal{T}_{g,m}$ and $v \in T_{x_0}\mathcal{T}_{g,m}$. Let $h\colon \mathbb{D} \to \mathcal{T}_{g,m}$ be a holomorphic mapping with $h(0) = x_0$ and $h_*((\partial/\partial\lambda)\mid_{\lambda=0}) = v$. Set $E^1(\lambda) = \mathrm{Ext}_{h(\lambda)}(F_1)$, $E^2(\lambda) = \mathrm{Ext}_{h(\lambda)}(F_2)$ and $E = a_1E^1 + a_2E^2$ for simplicity.

Then, the Levi form of ρ is

$$\mathcal{L}(\rho)[v, \overline{v}] = \frac{c\, E_{\lambda\overline{\lambda}}|_{\lambda=0}}{(c + E(0))^3} + \frac{E(0)\, E_{\lambda\overline{\lambda}}|_{\lambda=0} - 2|\, E_\lambda|_{\lambda=0}\, |^2}{(c + E(0))^3}.$$

Therefore, we conclude from Corollary 5.1 that

$$\begin{aligned}
&E(0)\ E_{\lambda\overline{\lambda}}\big|_{\lambda=0} - 2|\ E_{\lambda}|_{\lambda=0}\,|^2 \\
&= a_1^2 \left(E^1(0)\ E^1_{\lambda\overline{\lambda}}\big|_{\lambda=0} - 2|\ E^1_{\lambda}\big|_{\lambda=0}\,|^2\right) + a_2^2 \left(E^2(0)\ E^2_{\lambda\overline{\lambda}}\big|_{\lambda=0} - 2|\ E^2_{\lambda}\big|_{\lambda=0}\,|^2\right) \\
&\quad + a_1a_2 \left(E^1(0)\ E^2_{\lambda\overline{\lambda}}\big|_{\lambda=0} + E^2(0)\ E^1_{\lambda\overline{\lambda}}\big|_{\lambda=0} - 4\mathrm{Re}\left(\overline{E^1_{\lambda}\big|_{\lambda=0}}\ E^2_{\lambda}\big|_{\lambda=0}\right)\right) \\
&\geq a_1a_2 \left(E^1(0)\ E^2_{\lambda\overline{\lambda}}\big|_{\lambda=0} + E^2(0)\ E^1_{\lambda\overline{\lambda}}\big|_{\lambda=0} - 4\left|E^1_{\lambda}\big|_{\lambda=0}\right|\left|E^2_{\lambda}\big|_{\lambda=0}\right|\right) \\
&\geq a_1a_2 \Bigg(E^1(0)\ E^2_{\lambda\overline{\lambda}}\big|_{\lambda=0} + E^2(0)\ E^1_{\lambda\overline{\lambda}}\big|_{\lambda=0} \\
&\qquad -2E^1(0)^{1/2}\left(E^2_{\lambda\overline{\lambda}}\big|_{\lambda=0}\right)^{1/2} E^1(0)^{1/2}\left(E^2_{\lambda\overline{\lambda}}\big|_{\lambda=0}\right)^{1/2}\Bigg) \\
&= a_1a_2 \left(E^1(0)^{1/2}\left(E^2_{\lambda\overline{\lambda}}\big|_{\lambda=0}\right)^{1/2} - E^1(0)^{1/2}\left(E^2_{\lambda\overline{\lambda}}\big|_{\lambda=0}\right)^{1/2}\right)^2 \geq 0
\end{aligned}$$

and hence

$$\mathcal{L}(\rho)[v,\overline{v}] \geq \frac{c\ E_{\lambda\overline{\lambda}}\big|_{\lambda=0}}{(c+E(0))^3} = c\frac{a_1^2\mathcal{L}(\mathrm{Ext.}(F_1))[v,\overline{v}] + a_2^2\mathcal{L}(\mathrm{Ext.}(F_2))[v,\overline{v}]}{(c + a_1\mathrm{Ext}_{x_0}(F_1) + a_2\mathrm{Ext}_{x_0}(F_2))^3}$$

when $a+b>0$, $c>0$. Indeed, since $q_{F,x}$ is generic, the anti-complex linear map

$$T_{x_0}\mathcal{T}_{g,m} \ni v \mapsto \eta_v \in \mathcal{Q}_{x_0}$$

is an isomorphism. Therefore, ρ is (real analytic) strictly plurisubharmonic on $\mathcal{T}_{g,m}$. From the definition, ρ satisfies that $\rho(x) < 0$ for $x \in \mathcal{T}_{g,m}$. Since F_1 and F_2 are transverse, for any $\epsilon > 0$, there is a compact set $K \subset \mathcal{T}_{g,m}$ such that $\rho > -\epsilon$ for all $x \in \mathcal{T}_{g,m} - K$ (cf. §2.4). Since $\rho(x) > -1/a$ for all $x \in \mathcal{T}_{g,m}$, the function $\rho(x)$ satisfies the desired condition. □

We also obtain the following comparizon.

Theorem 5.3 (Gradients and Levi forms for log-extremal length functions)**.** *For any measured foliation F,*

$$d\log\mathrm{Ext.}(F) \wedge d^c\log\mathrm{Ext.}(F) \leq \frac{1}{2\mathrm{Ext.}(F)}dd^c\,\mathrm{Ext.}(F) \leq dd^c\log\mathrm{Ext.}(F)$$

holds in the sense of currents, where $d = \partial + \overline{\partial}$ and $d^c = i(\overline{\partial} - \partial)$. Especially, the log-extremal length function $\log\mathrm{Ext.}(F)$ *is a real analytic strictly plurisubharmonic function on $\mathcal{T}_{g,m}$, when F is essentially complete.*

Proof. By approximation, we may assume that F is essentially complete. Then, the inequalities follow from the combinations of (5.2) and Corollary 5.1. □

6. GEMETRIC INTERPRETATION OF η_v

6.1. **Motivation.** Let $x_0 = (M_0, f_0) \in \mathcal{T}_{g,m}$. Suppose $q_0 = q_{F,x_0}$ is generic. Let $v = [\dot{\mu}] \in T_{x_0}\mathcal{T}_{g,m}$. We defined the quadratic differential $\eta_v \in \mathcal{Q}_{g,m}$ to satisfy

$$\langle v, \psi\rangle = \int_{M_0} \frac{\overline{\eta_v}}{|q_0|}\psi \tag{6.1}$$

for all $\psi \in \mathcal{Q}_{g,m}$ in Lemma 4.1 and Theorem 5.1. The definition of η_v implies as if the infinitesimal Beltrami differential $\dot{\mu}$ were infinitesimally equivalent to the "infinitesimal deformation" along $\overline{\eta_v}/|q_0|$ on M_{q_0} (cf. [14]). However, $\overline{\eta_v}/|q_0|$ is not



FIGURE 2. A schematic of the map $\mathcal{J}_{x_0}$.

a Beltrami differential in general, becauce it may have $1/|z|$-singularities at zeroes of q_0 and hence it could happen $\|\overline{\eta_v}/|q_0|\|_\infty = \infty$. Therefore, the above discussion does not interpret correctly the geometry of the anti-complex linear map

$$T_{x_0}\mathcal{T}_{g,m} \ni v \mapsto \eta_v \in \mathcal{Q}_{x_0} \tag{6.2}$$

in general. In this section, we shall try to give a geometric description of the correspondence (6.2).

6.2. **Geometric interpretation.** From [6, Theorem 5.8] and [10], the map

$$\mathcal{Q}_{g,m} \ni q \mapsto (\mathbf{h}(q), \mathbf{v}(q)) \in \mathcal{MF} \times \mathcal{MF} \tag{6.3}$$

is a real analytic diffeomorphism around q_0. Hence, the map

$$\mathcal{J} \colon \mathcal{T}_{g,m} \times \mathcal{T}_{g,m} \ni (x, y) \mapsto -q_{\mathbf{h}(q_{F,x}),y} \in \mathcal{Q}_{g,m} \tag{6.4}$$

is also a real analytic diffeomorphism on a neighborhood $N_1 \times N_1$ of (x_0, x_0) onto the image, where N_1 is a neighborhood of $x_0 = \pi(q_0) \in \mathcal{T}_{g,m}$. Notice that $\mathbf{h}(\mathcal{J}(x, y)) = \mathbf{h}(q_{F,x})$, $\mathcal{J}(x, y) \in \mathcal{Q}_y$ and $\mathcal{J}(x, x) = q_{F,x}$ for all $x, y \in N_1$. Let $\mathcal{J}_y(x) = \mathcal{J}(x, y)$ for $(x, y) \in N_1 \times N_1$ (Figure 2).

Theorem 6.1 (Geometric interpretation). *For $v \in T_{x_0}\mathcal{T}_{g,m}$, we have*

$$(\mathcal{J}_{x_0})_*[v] = -4\eta_v,$$

where $\eta_v \in \mathcal{Q}_{x_0}$ is taken as (6.1).

Proof. Consider the triangulation Δ_q and the train track τ_{q_0} on M_q for the horizontal foliation $|\mathrm{Im}(\sqrt{q})|$ of q (cf. [6, Theorem 5.8]). We may assume that Δ_q contains no horizontal edge (cf. [6, Lemma 5.7]). Consider the lifts $\tilde{\Delta}_q$ and $\tilde{\tau}_q$ of Δ_q and τ_q to $\tilde{M}_q$, respectivey. We orient each edge of $\tilde{\Delta}_q$ so that $\mathrm{Im}(\omega_q)$ is positive along the edge. Each branch b of $\tilde{\tau}_q$ intersects a unique edge e_b of $\tilde{\Delta}_q$. We orient b so that $b \cdot e_b = +1$, where $\cdot$ means the algebraic intersection number. By taking U sufficiently small, we may also assume that the marking $\tilde{M}_{q_0} \to \tilde{M}_q$ induces isomorphisms from $\tilde{\Delta}_{q_0}$ and $\tilde{\tau}_{q_0}$ to $\tilde{\Delta}_q$ and $\tilde{\tau}_q$, respectively.

From [6, Theorem 5.8], the tangential map of the map $\mathbf{h}\colon \mathcal{Q}_{x_0} \to \mathcal{MF}$ of the horizontal foliations satisfies

$$\mathrm{Im}\int_{M_{q_0}} \frac{\psi_1\overline{\psi_2}}{4|q_0|} = \omega_{Th}\left(\mathbf{h}_*(\psi_1)\,,\, \mathbf{h}_*(\psi_2)\right)_{\tau_{q_0}} \tag{6.5}$$

for $\psi_1, \psi_2 \in T_{q_0}\mathcal{Q}_{x_0} \cong \mathcal{Q}_{x_0}$, where $\omega_{Th}\left(\cdot\,,\,\cdot\right)_{\tau_{q_0}}$ is the Thurston's symplectic form on $W(\tau_{q_0})$, where $W(\tau_{q_0})$ is the space of real valued functions on the branches of τ_{q_0} satisfying the switch condition. It is known that the space $W(\tau_{q_0})$ is identified with the tangent space of $\mathcal{MF}$ at $\mathbf{h}(q_0)$ since τ_{q_0} is maximal (cf. [29, §3.2]).

From the definition, $|\mathrm{Im}(\omega_{q_{F,x}})|$ is the lift of the horizontal foliation of $q_{F,x}$ to $\tilde{M}_{q_{F,x}}$. By the definition of the orientation of edges of $\tilde{\Delta}_{q_0}$, the image of the map $N_1 \ni x \mapsto \mathbf{h}(q_{F,x}) \in \mathcal{MF}$ at x is identified with $w_x \in W(\tilde{\tau}_{q_0})$ which is defined by

$$w_x(b) = \int_b |\mathrm{Im}(\omega_{q_{F,x}})| = \int_b \mathrm{Im}(\omega_{q_{F,x}}).$$

From Lemmas 3.1 and 4.1, the image of $v \in T_{x_0}\mathcal{T}_{g,m}$ under the differential of the map $N_1 \ni x \mapsto \mathrm{Im}(\chi_{q_{F,x}})$ is identified with the homomorphism

$$V_v\colon H_1(\tilde{M}_{q_0}) \ni c \mapsto -2\mathrm{Im}\int_c \frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}} \in \mathbb{R}. \tag{6.6}$$

Indeed, when $\{q_\lambda\}_\lambda \in D$ is taken as Lemma 4.1, (6.6) is derived from

$$\begin{aligned}
\frac{\partial \mathrm{Im}(\chi_{q_\lambda}(c))}{\partial t} &= -i\left(\frac{\partial \chi_{q_\lambda}(c)}{\partial \lambda} + \frac{\partial \chi_{q_\lambda}(c)}{\partial \overline{\lambda}}\right) \\
&= -i\int_c \overline{\left(\frac{(\pi_{q_\lambda})^*(\eta_\lambda)}{\omega_{q_\lambda}}\right)} - \frac{(\pi_{q_\lambda})^*(\eta_\lambda)}{\omega_{q_\lambda}} \\
&= -2\mathrm{Im}\int_c \frac{(\pi_{q_\lambda})^*(\eta_\lambda)}{\omega_{q_\lambda}}
\end{aligned}$$

where $\lambda = t + is$, since the real part of χ_{q_λ} is constant on D.

Let $p\colon \tilde{\tau}_{q_0} \to \tau_{q_0}$ be the induced covering. For a weight $w \in W(\tau_{q_0})$, we define $C_w \in H_1(\tilde{M}_{q_0}, \mathbb{R})^-$ as the homology class of a cycle

$$\sum_{b\text{:branches of } \tilde{\tau}_{q_0}} w(p(b))b.$$

Then,

$$\begin{aligned}
w_x(C_w) &= \sum_b w(p(b))w_x(b) \\
V_v(C_w) &= -2\mathrm{Im}\int_{C_w} \frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}} = -2\sum_b w(p(b))\mathrm{Im}\int_b \frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}}.
\end{aligned}$$

These equations imply that the differential of the map $N_1 \ni x \mapsto w_x \in W(\tilde{\tau}_{q_0})$ at $x = x_0$ is equal to the weight (which we abbreviate as)

$$V_v(b) = -2\,\mathrm{Im}\int_b \frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}}.$$

The weight V_v is invariant under the action of covering transformation r_0 of $\tilde{M}_{q_0} \to M_{q_0}$ on $\tilde{\tau}_{q_0}$, because for any branch b of $\tilde{\tau}_{q_0}$, the orientation of $r_0(b)$ is opposite to the orientation induced from b. Thus, V_v descends to a weight $p_*(V_v)$ on τ_{q_0}.

Let $\psi_0 = (\mathcal{J}_{x_0})_*[v]$. For all $\psi \in \mathcal{Q}_{x_0} \subset T_{q_0}\mathcal{Q}_{g,m}$, we define $w(\psi) \in W(\tilde{\tau}_{q_0})$ by

$$w(\psi)(b) = \mathrm{Im}\int_b \frac{(\pi_{q_0})^*(\psi)}{2\omega_{q_0}}.$$

Then, we have $\mathbf{h}_*[\psi] = p_*(w(\psi))$ (cf. [6, Proof of Theorem 5.8]). By the definition of $\mathcal{J}_{x_0}$, $\mathbf{h}\circ\mathcal{J}_{x_0}(x) = \mathbf{h}(q_{F,x})$ for all $x \in N_1$. Hence, $\mathbf{h}_*[\psi_0] = p_*(V_v)$ as elements of $W(\tilde{\tau}_{q_0}) \cong T_{\mathbf{h}(q_0)}\mathcal{MF}$. From (6.5) and [6, (5.4)], the following holds for all $\psi \in \mathcal{Q}_{x_0}$:

$$\begin{aligned}
\mathrm{Im}\int_{M_{q_0}} \frac{\psi_0\overline{\psi}}{4|q_0|} &= \frac{1}{2}\,\mathrm{Im}\int_{\tilde{M}_{q_0}} \frac{(\pi_{q_0})^*(\psi_0)\overline{(\pi_{q_0})^*(\psi)}}{4|\omega_{q_0}|^2}\\
&= \frac{1}{2}\,\omega_{Th}\left(V_v\,,\,w(\psi)\right)_{\tilde{\tau}_{q_0}}\\
&= -2\int_{\tilde{M}_{q_0}} \mathrm{Im}\frac{(\pi_{q_0})^*(\eta_v)}{2\omega_{q_0}} \wedge \mathrm{Im}\frac{(\pi_{q_0})^*(\psi)}{2\omega_{q_0}}\\
&= -\mathrm{Re}\int_{\tilde{M}_{q_0}} \frac{(\pi_{q_0})^*(\eta_v)}{2\omega_{q_0}} \wedge \overline{\frac{(\pi_{q_0})^*(\psi)}{2\omega_{q_0}}}\\
&= -2\mathrm{Im}\int_{\tilde{M}_{q_0}} \frac{(\pi_{q_0})^*(\eta_v)\overline{(\pi_{q_0})^*(\psi)}}{4|\omega_{q_0}|^2} = -\mathrm{Im}\int_{M_{q_0}} \frac{\eta_v\overline{\psi}}{|q_0|}.
\end{aligned}$$

Therefore, we have $(\mathcal{J}_{x_0})_*[v] = \psi_0 = -4\eta_v$. □

7. Appendix

7.1. **Alternative approach to Gardiner's formula.** Applying our method in Theorem 5.1, one can also get an alternative proof of Gardiner's formula for the derivative of extremal length in some case. Indeed, this is the case where we need in the proof of Theorem 1.1.

Let $F \in \mathcal{MF}$ and $x_0 = (M, f) \in \mathcal{T}_{g,m}$. Suppose that the Hubbard-Masur differential $q_0 = q_{F,x_0}$ is generic. Let $v = [\dot{\mu}]$ and $\{q_\lambda\}_{\lambda\in D}$ as the proof of Theorem 5.1. Suppose $x_\lambda = \pi(q_\lambda)$ varies holomorphically. By applying the discussion in the proof of Theorem 5.1, from Lemma 4.1, we have

$$\begin{aligned}
&\frac{\partial}{\partial\lambda}\mathrm{Ext}_{x_\lambda}(F)\Big|_{\lambda=0}\\
&= \frac{i}{4}\sum_{k=1}^{\tilde{g}}\left(\int_{\alpha_k}\overline{\left(\frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}}\right)}\int_{\beta_k}\overline{\omega_{q_0}} - \int_{\beta_k}\overline{\left(\frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}}\right)}\int_{\alpha_k}\overline{\omega_{q_0}}\right)\\
&\quad + \frac{i}{4}\sum_{k=1}^{\tilde{g}}\left(\int_{\alpha_k}\omega_{q_0}\int_{\beta_k}\overline{\left(-\frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}}\right)} - \int_{\beta_k}\omega_{q_0}\int_{\alpha_k}\overline{\left(-\frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}}\right)}\right)\\
&= \frac{i}{4}\int_{\tilde{M}_{q_0}}\overline{\left(\frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}}\right)}\wedge\overline{\omega_{q_0}} - \frac{i}{4}\int_{\tilde{M}_{q_0}}\omega_{q_0}\wedge\overline{\left(\frac{(\pi_{q_0})^*(\eta_v)}{\omega_{q_0}}\right)}\\
&= -\frac{1}{2}\int_{\tilde{M}_{q_0}}\frac{\overline{(\pi_{q_0})^*(\eta_v)}}{|\omega_{q_0}|^2}\omega_{q_0}^2 = -\int_{M_{q_0}}\frac{\overline{\eta_v}}{|q_0|}q_0 = -\int_M \dot{\mu}\,q_{F,x_0}
\end{aligned}$$

because $M = M_{q_0}$ by definition.

7.2. **Examples.** In this section, we treat the exceptional cases in the proofs of pluriharmonicity in the previous sections.

7.2.1. *Levi forms for the case of flat tori.* We start to consider the Levi forms of extremal length functions on the Teichmüller space of flat tori (once punctured tori), and check our formula in Theorem 5.1 also valid in this case. This case is excluded from our assumption in the theorem. However, this is a motivated example of our result.

The Teichmüller space of a flat torus is identified with the upper half plane $\mathbb{H}$. A point $\tau \in \mathbb{H}$ corresponds to a marked Riemann surface M_τ which is defined as the quotient space of $\mathbb{C}$ by a lattice generated by $z+1$ and $z+\tau$. Notice that the space $\mathcal{PMF} = \mathcal{PMF}(M_i)$ of projective measured foliations on M_i ($i = \sqrt{-1} \in \mathbb{H}$) is identified with the projective space $\mathbb{RP}^1 = \{[x\colon y] \mid (x,y) \neq (0,0)\}$. A point $[x\colon y] \in \mathbb{RP}^1$ corresponds to a measured foliation whose leaves are parallel to the line of direction $x+yi$.

Let α be a closed curve on M_i corresponding to $[1\colon 0] \in \mathcal{PMF}$. Then, the Jenkins-Strebel differential $q_{\alpha,\tau}$ on M_τ for α and the extremal length function of α at M_τ are obtained as

$$\begin{aligned} q_{\alpha,\tau} &= -\frac{1}{\mathrm{Im}(\tau)^2} dz^2 \\ \mathrm{Ext}_\tau(\alpha) &= \frac{1}{\mathrm{Im}(\tau)}. \end{aligned}$$

Fix $\tau_0 \in \mathbb{H}$. For $V \in \mathbb{C}$, let v be the tangent vector at τ_0 corresponding to the infinitesimal quasiconformal deformation from M_{τ_0} to $M_{\tau_0+\lambda V}$ as $\lambda \to 0$. The Levi form (Laplacian) of the extremal length function is

$$\mathcal{L}(\mathrm{Ext}.(\alpha))[v,\overline{v}] = \left.\frac{\partial^2}{\partial\lambda\partial\overline{\lambda}}\mathrm{Ext}_{\tau_0+\lambda V}(\lambda)\right|_{\lambda=0} = \frac{|V|^2}{2\mathrm{Im}(\tau_0)^3} \tag{7.1}$$

The Beltrami differential $\mu(\lambda)$ of the quasiconformal deformation from M_{τ_0} to $M_{\tau_0+\lambda V}$ behaves

$$\mu(\lambda) = \lambda\cdot\dot{\mu} + o(|\lambda|) = \lambda\cdot\frac{iV}{2\mathrm{Im}(\tau_0)}\frac{d\overline{z}}{dz} + o(|\lambda|)$$

as $t \to 0$. By definition, $v = [\dot{\mu}]$. We define a holomorphic quadratic differential η_v on M_{τ_0} by

$$\eta_v = -\frac{i\overline{V}}{2\mathrm{Im}(\tau_0)^3} dz^2. \tag{7.2}$$

This differential satisfies that

$$\langle v, \psi\rangle = \int_{M_{\tau_0}} \dot{\mu}\psi = \int_{M_{\tau_0}} \frac{\overline{\eta_v}}{|q_{\alpha,\tau_0}|}\psi$$

for all holomorphic quadratic differential ψ on M_{τ_0}. Therefore, the right-hand side of the formula in Theorem 5.1 is equal to

$$2\int_{M_{\tau_0}} \frac{|\eta_v|^2}{|q_{\alpha,\tau_0}|} = \frac{|V|^2}{2\mathrm{Im}(\tau_0)^3},$$

which coincides with the right-hand side of (7.1).

Let F be a measured foliation corresponding to $[a\colon b] \in \mathcal{PMF}$ with $b \neq 0$. We normalize F with $i(\alpha, F) = 1$. Then the Hubbard-Masur differential $q_{F,\tau}$ on M_τ

and the extremal length function for F are

$$q_{F,\tau} = -\frac{(a+b\overline{\tau})^2}{b^2\mathrm{Im}(\tau)^2}dz^2$$

$$\mathrm{Ext}_\tau(F) = \frac{|a+b\tau|^2}{b^2\mathrm{Im}(\tau)}.$$

Let v be the tangent vector at τ_0 corresponding to $\dot{\mu} = \frac{iV}{2\mathrm{Im}(\tau_0)}\frac{d\overline{z}}{dz}$. Then,

$$\mathcal{L}(\mathrm{Ext}_{\cdot}(\alpha))[v,\overline{v}] = \frac{\partial^2}{\partial\lambda\partial\overline{\lambda}}\mathrm{Ext}_{\tau_0+\lambda V}(\lambda)\Big|_{\lambda=0} = \frac{|a+b\tau_0|^2}{2b^2\mathrm{Im}(\tau_0)^3}|V|^2 \tag{7.3}$$

In this case, the differential $\eta_v \in \mathcal{Q}_{\tau_0}$ is

$$\eta_v = -\frac{i|a+b\tau_0|^2\overline{V}}{2b^2\mathrm{Im}(\tau_0)^3}dz^2. \tag{7.4}$$

and hence

$$2\int_{M_{\tau_0}}\frac{|\eta_v|^2}{|q_{F,\tau_0}|} = \frac{|a+b\tau_0|^2}{2b^2\mathrm{Im}(\tau_0)^3}|V|^2,$$

which coincides with the right-hand side of (7.3).

7.2.2. *The gemetric interpretation of η_v for the case of tori.* We check Theorem 6.1 in the case of flat tori (once punctured tori). We use the notation defined in §7.2.1 frequently.

First we consider the case $\alpha = [1\colon 0]$. Fix $\tau_0 \in \mathbb{H}$. Since the projective class of the horizontal foliation of $q_{\alpha,\tau}$ corresponds to a projective class $[-\mathrm{Re}(\tau)\colon 1] \in \mathbb{RP}^1$, the underlying foliation of the horizontal foliation $J_{\tau_0}(\tau)$ is foliated by the lines (leaves) directed to $-(\mathrm{Re}(\tau)) + \tau_0$. Since the horizontal foliations of $J_{\tau_0}(\tau)$ and $q_{\alpha,\tau}$ are same,

$$\int_\alpha |\mathrm{Im}\sqrt{J_{\tau_0}(\tau)}| = i(\mathbf{h}(J_{\tau_0}(\tau)),\alpha) = i(\mathbf{h}(q_{\alpha,\tau}),\alpha) = \int_\alpha |\mathrm{Im}\sqrt{q_{\alpha,\tau}}|$$

holds. Therefore, we can check that

$$J_{\tau_0}(\tau) = \left(\frac{-\mathrm{Re}(\tau)+\overline{\tau_0}}{\mathrm{Im}(\tau)\mathrm{Im}(\tau_0)}\right)^2 dz^2 \in \mathcal{Q}_{\tau_0}$$

for $\tau \in \mathbb{H}$. Let $v = [\dot{\mu}]$ be a tangent vector at τ_0, where $\dot{\mu} = \frac{iV}{2\mathrm{Im}(\tau_0)}\frac{d\overline{z}}{dz}$ with $V \in \mathbb{C}$ as §7.2.1. Then, from the above calculation, we obtain

$$(J_{\tau_0})_*[v] = \left(2\frac{i\overline{V}}{\mathrm{Im}(\tau_0)^3}\right)dz^2 \in \mathcal{Q}_{\tau_0}.$$

Comparing with (7.2), we get $(J_{\tau_0})_*[v] = -4\eta_v$ as we appeared in Theorem 6.1.

Let $F \in \mathcal{PMF}$ be a measured foliation corresponding to $[a\colon b] \in \mathcal{PMF}$ with $b \neq 0$ and $i(\alpha, F) = 1$. Then, the underlying foliation of the horizontal foliation of $q_{F,\tau}$ is foliated by the lines of direction $[a\mathrm{Re}(\tau) + b|\tau|^2\colon a + b\mathrm{Re}(\tau)]$. By the similar argument as above, one can check that

$$J_{\tau_0}(\tau) = \left(\frac{-(a\mathrm{Re}(\tau)+b|\tau|^2) + (a+b\mathrm{Re}(\tau))\overline{\tau_0}}{b\mathrm{Im}(\tau)\mathrm{Im}(\tau_0)}\right)^2 dz^2$$

$$(J_{\tau_0})_*[v] = \left(2\frac{i|a+b\tau_0|^2\overline{V}}{b^2\mathrm{Im}(\tau_0)^3}\right)dz^2 \in \mathcal{Q}_{\tau_0}.$$

From (7.4), we also have $(J_{\tau_0})_*[v] = -4\eta_v$.

7.2.3. *A comment on the case of* $3g-3+m=1$. The case of once punctured tori is treated in the same way as that in the case of flat tori. Indeed, the canonical completion at the puncture gives an isomorphism between the Teichmüller space of once punctured tori and that of flat tori. The extremal length functions in the both spaces are identified under the isomorphism. The case of fours punctured sphere is treated in the similar way.

References

[1] Gunnar Berg. Hyperconvexity and the Carathéodory metric. *Arch. Math. (Basel)*, 32(2):189–191, 1979.

[2] Lipman Bers and Leon Ehrenpreis. Holomorphic convexity of Teichmüller spaces. *Bull. Amer. Math. Soc.*, 70:761–764, 1964.

[3] Francis Bonahon. Bouts des variétés hyperboliques de dimension 3. *Ann. of Math. (2)*, 124(1):71–158, 1986.

[4] A. Douady and J. Hubbard. On the density of Strebel differentials. *Invent. Math.*, 30(2):175–179, 1975.

[5] Fathi A. Fried F. Laudenbach F. Ponaru V. Douady, A. and M. Shub. *Travaux de Thurston sur les surfaces*, volume 66 of *Astérisque*. Séminaire Orsay.

[6] D. Dumas. Skinning maps are finite-to-one. *ArXiv e-prints*, March 2012.

[7] Clifford J. Earle. On the Carathéodory metric in Teichmüller spaces. In *Discontinuous groups and Riemann surfaces (Proc. Conf., Univ. Maryland, College Park, Md., 1973)*, pages 99–103. Ann. of Math. Studies, No. 79. Princeton Univ. Press, Princeton, N.J., 1974.

[8] H. M. Farkas and I. Kra. *Riemann surfaces*, volume 71 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1992.

[9] Frederick P. Gardiner. *Teichmüller theory and quadratic differentials*. Pure and Applied Mathematics (New York). John Wiley & Sons, Inc., New York, 1987. A Wiley-Interscience Publication.

[10] Frederick P. Gardiner and Howard Masur. Extremal length geometry of Teichmüller space. *Complex Variables Theory Appl.*, 16(2-3):209–237, 1991.

[11] Roger Godement. *Topologie algébrique et théorie des faisceaux*. Actualit'es Sci. Ind. No. 1252. Publ. Math. Univ. Strasbourg. No. 13. Hermann, Paris, 1958.

[12] Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*. Wiley-Interscience [John Wiley & Sons], New York, 1978. Pure and Applied Mathematics.

[13] John Hubbard and Howard Masur. Quadratic differentials and foliations. *Acta Math.*, 142(3-4):221–274, 1979.

[14] Y. Imayoshi and M. Taniguchi. *An introduction to Teichmüller spaces*. Springer-Verlag, Tokyo, 1992.

[15] Yujiro Kawamata. On deformations of compactifiable complex manifolds. *Math. Ann.*, 235(3):247–265, 1978.

[16] Steven P. Kerckhoff. The asymptotic geometry of Teichmüller space. *Topology*, 19(1):23–41, 1980.

[17] Maciej Klimek. *Pluripotential theory*, volume 6 of *London Mathematical Society Monographs. New Series*. The Clarendon Press, Oxford University Press, New York, 1991. Oxford Science Publications.

[18] Kunihiko Kodaira. *Complex manifolds and deformation of complex structures*, volume 283 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1986. Translated from the Japanese by Kazuo Akao, With an appendix by Daisuke Fujiwara.

[19] S. L. Krushkal′. Strengthening pseudoconvexity of finite-dimensional Teichmüller spaces. *Math. Ann.*, 290(4):681–687, 1991.

[20] Samuel L. Krushkal. The Green function of Teichmüller spaces with applications. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):143–147, 1992.

[21] P. Lelong. *Fonctions plurisousharmoniques et formes différentielles positives*. Gordon & Breach, Paris-London-New York (Distributed by Dunod éditeur, Paris), 1968.

[22] Anna Lenzhen and Kasra Rafi. Length of a curve is quasi-convex along a Teichmüller geodesic. *J. Differential Geom.*, 88(2):267–295, 2011.

[23] L. Liu and W. Su. Variations of extremal length function on Teichmuller space. *ArXiv e-prints*, October 2012.

[24] Howard Masur. The Teichmüller flow is Hamiltonian. *Proc. Amer. Math. Soc.*, 123(12):3739–3747, 1995.

[25] Howard Masur and John Smillie. Hausdorff dimension of sets of nonergodic measured foliations. *Ann. of Math. (2)*, 134(3):455–543, 1991.

[26] Curtis T. McMullen. Complex earthquakes and Teichmüller theory. *J. Amer. Math. Soc.*, 11(2):283–320, 1998.

[27] Yair N. Minsky. Teichmüller geodesics and ends of hyperbolic 3-manifolds. *Topology*, 32(3):625–647, 1993.

[28] Takeo Ohsawa. On the Bergman kernel of hyperconvex domains. *Nagoya Math. J.*, 129:43–52, 1993.

[29] R. C. Penner and J. L. Harer. *Combinatorics of train tracks*, volume 125 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1992.

[30] Tibor Radó. *On the problem of Plateau. Subharmonic functions*. Springer-Verlag, New York-Heidelberg, 1971. Reprint.

[31] Hiroshige Shiga. On analytic and geometric properties of Teichmüller spaces. *J. Math. Kyoto Univ.*, 24(3):441–452, 1984.

[32] Jean-Luc Stehlé. Fonctions plurisousharmoniques et convexité holomorphe de certains fibrés analytiques. *C. R. Acad. Sci. Paris Sér. A*, 279:235–238, 1974.

[33] William Thurston. *The Geometry and Topology of Three-Manifolds*. Lecture Note at Princeton University, 1980. Available at `http://library.msri.org/nonmsri/gt3m/`.

[34] William A. Veech. The Teichmüller geodesic flow. *Ann. of Math. (2)*, 124(3):441–530, 1986.

[35] William A. Veech. Moduli spaces of quadratic differentials. *J. Analyse Math.*, 55:117–171, 1990.

[36] Scott A. Wolpert. Geodesic length functions and the Nielsen problem. *J. Differential Geom.*, 25(2):275–296, 1987.

Department of Mathematics, Graduate School of Science, Osaka University, Machikaneyama 1-1, Toyonaka, Osaka 560-0043, Japan